


Royal University of Phnom Penh Department of Psychology



Cambodian Mental Health Survey 2012



Dr. Tanja Schunert (PhD Psychiatry and Psychotherapy, GIZ, Germany)

Sareth Khann (MA Psychology, Department of Psychology, Royal University of Phnom Penh, Cambodia)

Sovandara Kao (BA Psychology & MA Psychology Candidate, Department of Psychology, Royal University of Phnom Penh, Cambodia)

Channika Pot (BA Psychology, Department of Psychology, Royal University of Phnom Penh, Cambodia)

Laura Bebra Saupe (BA Psychology, GIZ, Germany)

Sisokhom Sek (MA Psychology, Head of Department of Psychology, Royal University of Phnom Penh, Cambodia),

Hema Nhong (MA Psychology, Vice Dean of Faculty of Social Sciences and Humanities, Royal University of Phnom Penh, Cambodia)



This research was funded by GIZ (Gesellschaft für internationale Zusammenarbeit), Maryknoll, and Catholic Relief Services (CRS). We are indebted to these generous donors for funding the preparation, data collection, analysis and report writing of this large project. A special thanks to Maryknoll missioners Sister Luise Ahrens and Father Kevin Conroy and to Elizabeth Högger for assistance in securing the funding and for moral support throughout the project.

We express our sincere thanks to Dr. Sothy Eng, Dr. Cindy J. Lahar and Dr. Rowena Beecham for valuable contributions of ideas to the structure and content of the survey and to Rowena Beecham also for assistance with the preparation of the data entry matrix. A special thanks and appreciation are in order to Dr. Bunnack Poch who provided invaluable statistical expertise while developing the biographic interview questions, the sampling methods and the analyses. We express our sincere thanks to KC and Dr. Cindy J. Lahar for extensive editing assistance of the final report.

The following students of the Department of Psychology assisted in collecting data: Punreay But, Chanthy Chen, Sophea Chen, Sony Chuon, Tey Heng, Naysim Hong, Kosal Ith, Sovandara Kao, Somony Keo, Vinhean Keo, Samnang Long, Sophea Ngeth, Channa Pil, Khemroth Pho, Channika Pot, Pearum San, Sreyleak Saosin, Sopheak Sar, Chankanha Sim, Saren Svay, Socheat Thong, Bunthorn Thorng, Sokhom Som, Sovon Yeng, Sothearoath Yi and Chanthorn You.

We are very thankful to those psychology graduate students who served as team leaders during the data collection: Bunna Peoun, Thearom Ret, Thida Kim, Chansotheara Mut and Sareth Khann. Students of the Department of Psychology assisted in data entry: Sovandara Kao, Channika Pot, Samnang Long, Channa Pil, Sophea Ngeth, Sopheak Sar and Chanthorn You.

We are also grateful to all the Cambodian citizens who shared their time by answering the interview questions in their homes.



Aut	IOIS				
Ack	owledgements	ii			
Tab	able of contents				
List	of figures	vi			
List	able of contents ists of figures ists of tables vibbreviations vibbreviations vibstract lecommendations & summary introduction I. Mental health in Cambodia II. The development of a nationwide mental health survey at the Department of Psychology (DP) at the Royal University Phnom Penh (RUPP) Specific objectives of the Nationwide Mental Health Survey III. Situational background of the present survey Resources in mental health Mental health policy in Cambodia Cultural explanations for and presentations of mental illnesses in Cambodia Treatment seeking behavior and treatment options for mental illnesses in Cambodia Research findings regarding the different issues addressed in this survey 1) Violence and mental health 2) PTSD 3) Cultural validity of the PTSD construct 4) Anxiety and depression 5) Tendency to suicide 6) Second generation trauma 7) Addictive disorders 8) Domestic violence 9) Schizophrenia II. Sample III. Instruments, scales and survey development				
Abb		vii			
Abs	ract	iii vi viii X Xiii 1 2 hology (DP) 4 4 4 7 8			
Rec	mmendations & summary	Xii			
Intr	duction	1			
	I. Mental health in Cambodia	2			
	II. The development of a nationwide mental health survey at the Department of Psychology (Di	P)			
	at the Royal University Phnom Penh (RUPP)	4			
	Specific objectives of the Nationwide Mental Health Survey	4			
	III. Situational background of the present survey	4			
	Resources in mental health	4			
	Mental health policy in Cambodia	7			
	Cultural explanations for and presentations of mental illnesses in Cambodia	8			
	Treatment seeking behavior and treatment options for mental illnesses in Cambodia	9			
	Research findings regarding the different issues addressed in this survey	10			
	1) Violence and mental health	11			
	2) PTSD	11			
	3) Cultural validity of the PTSD construct	12			
	4) Anxiety and depression	13			
	5) Tendency to suicide	13			
	6) Second generation trauma	14			
	7) Addictive disorders	14			
	8) Domestic violence	15			
	9) Schizophrenia	15			
Me					
	•				
	•				
	III. Data collection procedures				
	IV. Data entry				
	V. Data analysis	20			

Results		22
I.	Biographical background - respondent profiles (gender, age, position)	22
	Marital status	22
	Education	23
	Occupation	23
	Family members	23
	Economic situation	23
	Ownership of property and landholdings	24
II.	Screening questions	25
	Children problems	25
	1) School problems	25
	2) Aggressive behavior	26
	3) Other problems	26
	4) Detention of children	27
	5) Percentage of respondents' children suffering from different problems	28
	Traumatic events experienced	28
	Mental disorders of family members	28
	Irritability/ aggressiveness	29
	Suicidal tendency	29
	1) Suicide attempts	29
	2) Committed suicides	30
	Alcohol problems	30
	1) Alcohol problems of family members	30
	2) Alcohol consumption of respondents	30
	Drugs	31
	Gambling	32
	Cross-tables for `suicide attempts´, `in cage or chained´ and `alcohol problems´ with	
	demographic variables	32
	Correlations between suicide attempts, in cage or chained and alcohol problems with	
	demographic variables (age, years of education and economic situation)	33
III	. Traumatic experiences	35
	Traumatic experiences during Civil War	35
	Traumatic experiences after Civil War	36
	Other stressful experiences	36
	Traumatic events combined	37
	Correlations between the different traumatic/stressful experiences	37
IV	. Self-reporting symptom scales	39
	Harvard Trauma Questionnaire Symptoms	39
	Cambodian Addendum of Culturally Sensitive Items (now: C-SSI)	40
	The Hopkins Symptom Checklist – 25 for Anxiety and Depression (HSCL-25)	41
	1) Symptoms of anxiety	42
	2) Symptoms of depression	43
	V. Mental health and demographic factors	45

Gender	45
Age	45
Marital status	46
Location	46
Education	47
Economic situation	47
Correlations between different symptom scores and demographic factors	48
VI. Services	49
Service utilization, costs and knowledge	49
Needs for services	50
Summary & discussion	51
I. Screening questionnaire	51
Children problems	51
Traumatic events	51
Probable psychotic/ schizoaffective/ bipolar disorders	52
Suicidal tendency	53
Addictive disorders	54
II. Traumatic experiences	55
III. Symptom scales	57
Posttraumatic stress symptoms (HTQ)	57
Cambodian Symptom and Syndrome Inventory (C-SSI)	58
Anxiety and depression (HSCL-25)	58
IV. Symptom scales and demographics	59
V. Help seeking behavior	61
Needs	61
Limitations & conclusions	62
Conclusions	62
Appendix	64

References

68



Figures:

Fig. 1:	Household position of respondents in %	22
Fig. 2:	Marital status of respondents in %	22
Fig. 3:	Educational status of sespondents in %	23
Fig. 4:	Main occupation of respondents in %	23
Fig. 5:	Amount of land owned by the respondents in %	24
Fig. 6:	Percentage of households with respondents' and other children having problems at school	25
Fig. 7:	Percentage of households with respondents' and other children showing aggressive behavior	26
Fig. 8:	Percentage of households with respondents' and other children suffering from other problems	26
Fig. 9:	Percentage of households with respondents' or other children being brought to reeducation	
	camps	27
Fig.10	: Percentage of households with respondents' or other children which had contact with the	
	police, court or went to prison	27
Fig.11	: Percentage of respondents' children suffering from various problems	28
Fig.12	: Percentage of attempted suicides within a given time frame, cumulated percentages	29
Fig.13	: Percentage of respondents consuming alcohol in given time periods	30
Fig.14	: Percentage of respondents consuming the given amount of alcohol per occasion	31
Fig.15	: Traumatic events experienced or witnessed by respondents during Civil War	35
Fig.16	: Traumatic events experienced or witnessed by respondents after Civil War in %	36
Fig.17	: Other difficult events experienced by respondents in %	37
Fig.18	: HTQ-Symptoms experienced quite a bit or extremely by > 9% of respondents	39
Fig.19	: C-SSI items experienced extremely or quite a bit by > 15% of the respondents	41
Fig. 20): HSCL-25 symptoms of anxiety experienced extremely or quite a bit by > 10% of respondents	42
Fig. 2	: HSCL-25 symptoms of depression experienced extremely or quite a bit by > 10% of respondents	43
Fig. 22	2: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by sex	45
Fig. 23	3: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by age split	45
Fig. 24	H: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by marital status	346
Fig. 2	5: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by location	46
Fig. 26	8: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by education	
	in categories	47
Fig. 27	7: Mean total scores of C-SSI, HSCL-25 and HTQ-Symptoms by savings per person in the	
	respondents' household	47
Fig. 28	3: Percentage of yes answers to questions about services economics and knowledge	49
Fig. 29	e): Percentage of respondents who approached the mentioned sources for help	49



Tables:

Table 1:	Marital status and number of respondents by region of origin	22
Table 2:	Financial situation of the respondents relative to demographics and average monthly income	24
Table 3:	Percentage of respondents with different types of houses	24
Table 4:	Cumulative percentages of respondents who experienced school problems in certain time frames	25
Table 5:	Period in which respondents' and other children in the households exhibited aggressive behavior	26
Table 6:	Mean symptom scores related to suicide attempts	30
Table 7:	Cross-table for respondents alcohol problems, suicide attempts and being held captive in a cage	
	or on chains and sex and living area	32
Table 8:	Frequencies of most and least experienced traumatic events	35
Table 9:	Inter-correlations between various traumatic experiences	38
Table 10:	: M, SD and CI for HTQ scores and percentage of probable PTSD diagnosis	39
Table 11:	: M, SD and CI of C-SSI total, C-SSI syndromes, C-SSI symptoms	40
Table 12:	: M, SD and CI for HSCL-25 scores of anxiety	41
Table 13:	: M, SD and CI for HSCL-25 scores of depression	42
Table 14:	: Spearman correlations between mean symptom scale scores and demographic factors with	
	significance level	48
Table 15:	: Traumatic experiences during CW in comparison with other studies conducted in Cambodia	55
Table 16:	: Traumatic experiences after CW in comparison with other studies conducted in Cambodia	56
Table 17:	: Prevalence rates of anxiety and depression in comparison with other studies conducted in	
	Cambodia	58



AFESIP Agir Pour les Femmes En Situation Précaire

CBT Cognitive Behavioral Therapy

CCAMH Centre for Child and Adolescent Mental Health
CDC Center for Disease Control and Prevention
CIDI-Scale Composite International Diagnostic Interview
C-SSI Cambodian Symptom and Syndrome Inventory

CW Civil War

CWCC Cambodian Women's Crisis Center

DBT Dialectic Behavioral Therapy

DC-CAM Documentation Center of Cambodia

DP Department of Psychology

DSM III-R/IV-TR Diagnostic Statistic Manual 3rd edition revised/ 4th edition text revised

DV Domestic Violence

ECCC Extraordinary Chambers of the Courts of Cambodia
EMDR Eye Movement Desensitization and Reprocessing

FHI Family Health International
GBD Global Burden of Disease

GIZ Gesellschaft für Internationale Zusammenarbeit

HAP Human Assistance Program

HSCL-25 Hopkins Symptom Checklist 25 Items

HTQ Harvard Trauma Questionnaire

ICESCR International Covenant on Economic Social and Cultural Rights

IOM International Organization for Migration

KR Khmer Rouge

LICADHO Cambodian League for the Promotion and Defense of Human Rights

M Mean

MOH Ministry of Health

MOWA Ministry of Women's Affaires

N Number

NGO Non-Governmental Organization
NET Narrative Exposure Therapy

OPD Outpatient Psychiatric Department
PADV Project Against Domestic Violence

PPS Probability Proportional to Size

PSF Psicólogos sin Fronteras

PTSD Post Traumatic Stress Disorder
RUPP Royal University Phnom Penh

SD Standard Deviation

SSC Social Services Cambodia

TF-CBT Trauma-Focused Cognitive Behavioral Therapy

TPO Transcultural Psychosocial Organization
UNDP United Nations Development Programme

UNICEF United Nations Children's Fund

USA United States of America
WHO World Health Organization

YFP Youth for Peace

YRDP Youth Resource Development Program



This study reports on the first large-scale investigation of mental health in Cambodia. A randomized sample of 2690 adults from 9 provinces across Cambodia were interviewed about mental health and mental disorders, addictive and violent behaviors, suicidal tendency, traumatic events, and culturally influenced symptoms related to mental health. In addition, the Cambodian versions of the Harvard-Trauma Questionnaire (HTQ) and the Hopkins Symptoms Checklist (HSCL-25) were administered to all respondents.

Results indicate high rates of children's problems, especially aggressive behavior in the families (11.5%) (for comparison in India a rate of 12.8% was found for child and adolescent disorders), child abuse experienced by males (10.6%) and by females (8%) (as compared to 10.2% of cases of maltreated children in the US), suicide (0.8% of the households, 42.35 suicides per 100,000 per year 2011 compared to worldwide numbers with an average of 16 suicides per 100,000 population per year (WHO, 2011a) and a Thai rate of less than one fifth of the Cambodian rate) and suicide attempts (115, 4.3% total, 1.7% male, 5.5% female). Probable schizophrenic disorders were estimated at 0.6% for male and 0.2% for female (based on individuals being chained or put in a cage).

The HTQ-Symptom scale results found 3.1% of women and 1.6% of men suffering from probable PTSD. From the HSCL-25, estimates were that 31.7% of women and 18.4% of men suffer from probable anxiety disorder and 19.7% of women and 10.2% of men suffer from probable depressive disorders. The most common depressive symptoms were worrying, difficulty sleeping and low energy.

Physical forms of domestic violence were committed twice as frequently by women as

compared to men, whereas financial and sexual forms of domestic violence was more frequently committed by men. The percentage of male respondents committing domestic violence under the influence of alcohol was about 6 times higher compared to females.

Of the respondents 64.1% had experienced the period of Civil War. Of these 92.3% had experienced at least one traumatic event, with a mean of 7.0. Traumatic events were also reported among those younger respondents who did not experience the Civil War, where 70.2% experienced at least one traumatic event (with a mean of 1.3). Most of these younger respondents (94.4%) had experienced one to 12 other stressful experiences such as death or severe illness of close family members, marital, problems or financial worries. In the full sample, 87.9% had experienced some traumatic event, 86% suffered from financial worries and 15.9% reported fear of landmines.

Scores on scales of depression, anxiety, and trauma showed significant correlations with demographic data variables such as age, sex and years of education. For example, traditionally culturally-influenced Cambodian symptoms were assessed by including a checklist of cultural sensitive items in a scale labeled the Cambodian Symptom and Syndrome Inventory (C-SSI). Scores on the C-SSI increased most with higher age and lower education. Gender-dependence was high for all of the scales with significantly higher proportions of women suffering from psychological symptoms. Increasing poverty is associated with increasing depressive symptoms, anxiety and trauma.

Only 24.1% of the respondents had ever sought help for the symptoms discussed in these

interviews. Some had approached a health centers (62.3%), some sought help within their family (50.7%) some at pharmacies (34.1%) and some contacted traditional, spiritual resources (about 25%). Nearly all (97.2%) wished for a mental health service at their local health center and 35.7% stated they would see a psychologist.

Major limitations of this study were that in approximately 50% of the interviews complete privacy was not obtained. Due to the planting season the randomized sampling led to an overrepresentation of female respondents (66%). Because the survey was conducted by interviews based on self-reports of respondents and no clinical/psychiatric examination took place, prevalence rates are estimations for probable disorders.

In summary a high percentage of mental disorders especially among the female population was found. Problems that need to be addressed with priority are suicidal behaviors, children and adolescent mental health, mental health and poverty issues as well as human rights, with an urgent need for improvement of access to treatment by psychiatrists and psychologists and of mental health literacy among the population.



Support and conduct further research in the fields of

- Etiological research (see also Stewart et al., 2010) especially on causes of suicidal behavior
- Second generation trauma
- Child and adolescent mental health
- Psychosis
- Epidemiology (Ministry of Health (MoH), 2011) with a need of data sharing
- Monitoring and evaluation of treatment interventions (see also Stewart, Tsong, Phan Chan, 2010)
- Further cross cultural studies (see also D. Hinton, A. Hinton, Eng & Choung, 2011)
- Bound reactions to traumatic events, culture dependent interpretation of psychological symptoms

On the governmental level

- Put the National Mental Health Plan into practice
- Increase nationwide services in the provinces: install governmental positions for psychiatrists and psychologists e.g. in hospitals and health care centers. Make such positions more attractive by increasing staff salaries (see also MoH, 2011, p. 10 and 19)
- Increase community mental health care with regular follow up care: cost effective, decentralized and services integrated into local settings in order to deliver an adequate coverage of mental health care for the population, including outpatient clinics, mobile community mental health teams for outreach, acute inpatient care, rehabilitation and occupation (see also MoH 2003; Saxena, Thornicroft, Knapp & Whiteford, 2007; Somasundarami & van de Put, 1999; WHO, 1990)
- Include multi professional teams that implement culturally appropriate, fitting interventions including psychiatrists, psychiatric nurses, psychologists, art therapists and social workers (see also MoH, 2011, p.21)
- Promote cooperation and a participatory approach among all stakeholders (see also MoH, 2011, p.13)
- Add positions for psychologists and counseling to psychiatric services and addiction programs in psychiatric hospitals

- Establish positions for school counselors: to promote positive mental health and mental health literacy in schools, to provide a valuable opportunity for better preparation for challenges in life, to help to prevent deterioration at an early stage, especially considering the high percentage of youth and adolescents in Cambodia (see also Stewart et al., 2010)
- Support parenting classes: to interrupt the cycle of intergenerational transmission of psychological problems, to promote positive mental health, communication and interaction skills
- Implement a mental health legislation for fair humane treatment: to protect the basic human and civil rights of people with mental disorders, especially those receiving involuntary treatments (MoH, 2003, Saxena,et al., 2007)
- Implement a health insurance system: to pool risks especially for the poor population as already in process by the Ministry of Labor and Vocational Training
- Establish a suicide prevention program: culturally appropriate and gender specific programs, combined with school educational programs on positive mental health (see also Jega Nathan & Kullgren 2011), stop public presentation of videos "glorifying" suicide
- Continue and amplify programs on domestic violence: to rise awareness in the population and offer assistance and support, to train more effective communication skills to families
- Focus on programs for prevention of depression and anxiety especially for women (WHO, 2011c)
- Promote mental health literacy (knowledge about signs and symptoms of mental disorders and whom to approach best) with psycho-education programs focusing on child raising, family relationships, and consequences of mental problems through/via mass media e.g. radio/TV, newspaper etc.
- Diminish poverty by creating more job opportunities/ income generating programs as a long term preventive measurement against mental disorders

For the Departments of Psychology, Social Work and training of psychiatrists

- Ensure high quality curricula
- Include client-based teaching approach, practical training in class and in governmental and NGO mental health services (see also MoH, 2011, p.11)
- Provide constant supervision for students' internships
- Ensure self-care and burnout prevention for staff and students
- Promote "mental health literacy" for the population
- Conduct further research on mental health issues
- Participate in international cooperation and collaboration

Promising counseling approaches

- Support and apply cross-cultural approaches utilizing local resources (see also Nickerson & Hinton, 2011; Boehnlein & Kinzie, 2011)
- Include family and group counseling
- Individualize treatment settings (Dubois et al., 2004): strength based, short term solution focused, CBT, client-centered (see also Stewart et al., 2010)

- Offer trauma focused counseling (EMDR, testimony and narrative therapies, self-help groups)
- Offer art therapeutic approaches for children
- Train elders as mediators, Buddhist monks and traditional healers (see also Stewart et al., 2010)
- Integrate meditation (e.g. breathing) mindfulness and awareness exercises from Buddhism into counseling and prevention programs
- Integrate Buddhist ceremonies in group counseling where appropriate
- Provide multi professional teams together with social workers to address the psychosocial needs, poverty reduction, vocational training
- Develop psychoeducation programs for psychosis, anxiety, depression, addictive disorders, aggressive behavior and other mental health problems

Introduction

"There is no health without mental health" said Ban Ki-moon on October 10th, 2011, the World Mental Health Day, "Mental disorders are major contributors to illness and premature death, and are responsible for 13% of the global disease burden" (Chinese Women's Research Network, 2011, Prince et al., 2007).

- Mental disorders exact a toll greater than tuberculosis, cancer, or heart disease (World Bank, 1993).
- Mental health is an integral and essential component of health as mental disorders are among the leading causes of disability (World Health Organization's - WHO 2005 estimation, Mathers & Loncar, 2005; Murray & Lopez, 1997).
- Worldwide epidemiological data show that prevalence rates for child and adolescent mental disorders are around 20%, and types of disorders vary only little across different cultures. It is alarming to realize that this data also suggest a very early start of psychological aberrant behaviour. About half of all lifetime mental disorders begin before the age of 14 years (Saxena, Thornicroft, Knapp & Whiteford, 2007).
- Mental illness unfortunately often develops into chronic disorders, especially if untreated for a long period of time due to a lack of services and economic resources or fear of losing face and feelings of shame. The average time that passes between first evidence of a disease and the initial more adequate treatment was estimated between 3 months

- and 7 years (Stewart, Tsong & Phan Chan, 2010).
- Regardless of these alarming facts, analysis of data from the WHO's Atlas project shows widespread, systematic, and long-term neglect of resources for mental health care in low and middle income countries (WHO, 2005, Saxena, Sharan, Garrido & Saraceno, 2006).
- This phenomenon is even more prominent in the poorest countries as they spend the smallest proportion of their already scarce resources on mental health (less than 1%) (Saxena et al., 2007).

Mental health worldwide

- No health without mental health
- Mental disorders account for about 13% of the global burden of disease
- Mental disorders are among the leading causes of disability
- The most vulnerable groups are the uneducated, female, young and rural people
- Long-term neglect of resources for mental health care in low- and middle-income countries
- The poorest countries spend the smallest proportion of their already scarce resources on mental health
- Poor provision of mental health is costly

The most vulnerable groups for mental health disorders are the indigent, uneducated, female,

young, and rural people (Saxena et al. 2007). Yet these groups have the lowest access to services. The WHO has reported that the treatment gap for serious mental disorders is 35 to 50% for developed countries and 76 to 85% for low and middle-income countries (WHO, 2004 a). The proportion of those receiving good quality and humane treatment is small or none. The consequent results of inadequate mental health care are exponential.

Consequent results of inadequate mental health care:

- Human rights violations
- Discrimination
- Individual suffering
- Denial of opportunities
- Violence

- Social exclusion
- Unemployment
- Disability
- Family disputes
- Economic loss

Indeed, mental health issues account for 3 to 4% of the Gross Domestic Product of developed countries. Costs for low-income countries are much higher due to high costs, financial impact on family caretakers, and losses in productivity (McDaid, Knapp & Raja, 2008).

Many governments and policymakers ignore the high cost of mental health issues due to the fact that it is less visible than physical disability and has a highly negative social stigma. Rejection and avoidance of mental health issues is a universal phenomenon (Saxena et al., 2007). In many cultures, the mentally ill are said to be possessed by evil spirits as a punishment for misbehaviour. Thus, many still believe treatment for mental disorders is not viable.

It will take a great deal of advocacy and effort to improve mental health literacy (i.e. the ability to correctly identify mental illness) and to change attitudes and perceptions. Additionally, human, social, and financial resources will have to be increased significantly in order to achieve the World Health Report objective of adequate access to effective and humane treatment for those who suffer from mental disorders (Saxena et al., 2007).

I Mental health in Cambodia

Cambodia is dealing with an exceptionally challenging and specific situation regarding mental health. The many decades of Civil War, brutal regimes, and intensive/prolonged persecution have few parallels in the twentieth century.

The Cambodian holocaust has left a deeply disturbed and traumatized society:

- The loss of over a quarter of the population (with an emphasize on Cambodian intelligentsia)
- Remaining landmines and unexploded bombs
- Destruction of infrastructure, health care, education systems, as well as religious and cultural legacies

The burden of having to restart and rebuild these entire human, economic, and societal structures has made Cambodia enormously dependent on external aid. The official development assistance was 884.5 million USD according to the Paris Declaration Survey 2011 (UNDP, 2010).

This transformation from a tranquil agrarian society into dehumanized brutality remains inexplicable. In survivor self-help groups conducted jointly by the Department of Psychology (DP) and the Trans-cultural Psychosocial Organization (TPO) the demand for explanation and the lack of understanding regarding how this could happen poses a major obstacle in the endeavour to find meaning and/or a potential healing process. This situation seems to amplify somatic symptoms found so commonly in Cambodia (see also: Perry, Oum, & Gray, 2007). The KR period has only been introduced in schools since 2009 by the Documentation Centre of Cambodia (DCCAM) and the Ministry of Education (Dy, 2009).

Mental health in Cambodia

- Loss of over a quarter of the population, legacy of destruction of culture and infrastructure
- Pure dehumanized brutality seems inexplicable, great demand for explanation and better understanding
- Vast and multiple traumatic experiences
- Great need for psychological support for participation at the ECCC of KR survivors
- Enormous dependence on external aid
- Promoting mental health care has the potential to relieve personal suffering, as well as to bring about significant benefits to the society at large

Concise list of atrocities

Between the early 1970s and late 1990s a single individual could experience (singly or in any combination) the following traumatic events:

- Displacement from Vietnam War
- Witnessing torture and/or murders •
- Separation from family
- Becoming a KR child soldier
- Betrayal of family members
- Killing others
- Poverty Humiliation
- Forced labour
- mily members No education/health care
- Refugee camps
- Bombings
- Land mines
- Loss of land
- Starvation

All of these experiences can contribute to personal multiple type 2- trauma (see also: Field, 2011)

In 2006 the "Khmer Rouge trials" (a hybrid tribunal sponsored jointly by the United Nations and the Kingdom of Cambodia) were established to try the primary KR leaders and formally known as the "Extraordinary Chambers in the Courts of Cambodia" (ECCC).

With the pending trials regarding case 002 at the ECCC, NGOs such as TPO are confronted with a number of no less than 3,866 recognized civil parties that urgently need psychological preparation before their participation during the hearings to prevent retraumatization.

National reconciliation, reconstruction, and development can only occur with the establishment of a mental health program to address motivation, self-esteem, friendly relationships, and self-confidence (Somasundaram & van de Put, 1999).

Barriers towards national reconciliation (Human Rights Watch, 2010; Humeniuk, Ali, & Ling, 2004; van de Put & Eisenbruch, 2002)

Socio-political issues:

•Violence •Poverty •Scarcity of services

•Lack of education/safety •Unemployment

•Repression • Malnutrition • Human rights

Interlinked with

Post-traumatic symptoms:

•Nightmares •Poor sleep •Hyper vigilance

•Loss of self-control •Mistrust •Fear

•Varied somatic symptoms •Substance abuse

•Poor concentration •Stigmatization

However, child abuse, sexual exploitation, domestic violence, trafficking, gambling, and alcohol dependence pose serious problems in Cambodia and lead to a further decrease in mental health (Somasundaram & van de Put, 1999). These issues have therefore been included in the screening part of the present survey's questionnaire.

II The Development of a Nationwide Mental Health Survey at the Department of Psychology (DP) at the Royal University Phnom Penh (RUPP)

A significant lack of epidemiological data, training, and services in mental health was jointly discussed by the DP on September 24th 2010 at the WHO office in Phnom Penh. As per WHO guidelines (WHO, 2004 b and 2008), the DP decided to explore the extent of psychological problems and mental disorders in Cambodia as well as possible etiological, historical and socioeconomic contributing and/or resulting factors.

Thus, the DP decided to conduct a nationwide survey on mental health to contribute to a more comprehensive and in-depth situational analysis. This epidemiological research included cultural treatment-seeking behaviour and will help to provide a informative baseline for future planning of interventions and policy in order to ensure efficiency and address service gaps (see also: Stewart et al., 2010).

Specific objectives of the Nationwide Mental Health Survey

The purpose of the survey is to provide information for program planning on the part of the DP, the Ministry of Health and other institutions working in the field of mental health.

Objectives of this survey

- Provide reliable population-based information/figures about important mental-health-related issues such as violence, addictive behaviors and psychological symptoms in order to identify positive and negative trends and thus target efforts in an evidenced-based way
- Provide reliable population-based information about mental health service utilization and needs for planning purposes
- Assess major topics and needs related to mental health in order to help adjust programs and offers to the Cambodian population
- Establish a baseline against which program long-term effectiveness and impact can be subsequently assessed
- Build capacity in the area of research, monitoring and evaluation at the DP
- Establish more cooperation between the DP and the Ministry of Health
- Advocate and raise awareness for psychological issues and mental health

III Situational background of the present survey

Resources in mental health



Most of the staff and patients of the 800-bed psychiatric hospital in Takhmau were killed during the KR regime. In 1989, the United Nations Development Program estimated that no more than 300 qualified persons of all disciplines were available to serve the country (Bit, 1991). In addition, Buddhist principles foster an acceptance of suffering. Cambodian culture does not support socially acceptable ways of expressing and releasing anger/frustration (Bit, 1991).

Legacy of history

- During the Pol Pot era (1975-1979), it is estimated that approximately 2 million Cambodians (>1/4 of the population) died
- Only 50 physicians out of 1,000 (including two psychiatrists) before 1975 survived this period, with no mental health professionals among them (Stewart et al., 2010)
- No more than 300 qualified persons of all disciplines were left to serve the country (Bit, 1991)
- No conceptual framework or system of mental health care in Cambodia until 1993, trauma-related and other mental disorders largely went untreated (Hinton, Hinton, Eng & Choung, 2011a)

In 1994, the International Organization for Migration (IOM) and the University of Oslo, in cooperation with the Ministry of Health in Cambodia, initiated the training of psychiatrists. At present, 49 psychiatrists (11 females among them) have been trained (according to the University of Health Sciences). Additionally, 45 psychiatric nurses have been trained and work in mental health facilities and/or private practice in Cambodia (information from Ministry of Health). Mental health care has also been integrated into primary health care by the Ministry of Health and includes 297 trained physicians and 270 trained nurses.

However, according to several Cambodian psychiatrists, psychiatric education doesn't include extensive training in counselling and psychothera-

peutic skills, and the National Program for Mental Health estimates that 30% of psychiatrists and 90% of the trained primary care professionals are not involved in clinical mental health care (Stewart et al., 2010, Ministry of Health, 2010).

There currently exist 68 outpatient psychiatric departments (OPD) under the Ministry of Health. Of these, 50 exist among Cambodia's 84 referral hospitals and 18 are located among the 967 health centres (Ministry of Health, 2010). There are 8 OPDs at national level and one includes services for children at the Clinic for Child and Adolescent Mental Health (CCAMH) - Chey Chumnea provincial hospital in Takhmau.

In 2010, mental health services were available in 9 out of 24 provinces and 12,000 new psychiatric cases (32% male and 68% female) were diagnosed at governmental mental health services (Ministry of Health, 2010).

Nationwide, there are two psychiatric in-patient units at two referral hospitals with 16 acute care psychiatric beds in Phnom Penh (Khmer Soviet Friendship Hospital and Sunrise Clinic). However, there is no constant routine follow-up community care provided by the governmental health sector and only few NGOs provide community based care in the provinces. There were two psychosocial rehabilitation centres at Phnom Penh (1997) and Siem Reap (2001), but funding is very short and some programs had to be stopped.

The DP at RUPP has existed since 1994 and offers degree programs in psychology (undergraduate since 1994 and postgraduate since 2008). Additional degree courses are provided in social work (BA since 2008 and MA since 2009). However, no psychologists have yet been involved in governmental services such as hospital care or education. Regardless, the DP programs aim at meeting the great need for welltrained psychologists and counsellors in Cambodia's rural and urban areas. The vision includes setting up a cooperation network with a wider range of governmental and non-governmental institutions to establish synergies and enhance sustainability and efficiency.

Over 660 students have graduated since 1994 with a Bachelor's degree in psychology. Since 2008, the Department has offered a Master of Arts in Clinical Psychology and Trauma Treatment which provides advanced concepts and principles of psychology, psychotherapy, counselling, and the methods of scientific inquiry into the behaviour of individuals, groups and society. The goal is to train psychologists with a special focus on trauma therapeutic approaches along with a strong integration of traditional Cambodian cultural ways and forms of support.

NGOs often have a leading role in developing countries in the development of new services. However, they often supplement inadequate state infrastructure for mental health care as done by the Trans-cultural Psychosocial Organization (TPO) and Social Services Cambodia (SSC). Hence, poor collaboration across all sectors of different organizations and between practitioners of different professions (social work, psychiatry, and counselling) seems to worsen due to an increasing competition for limited funding (Reicherter, Boehnlein & Stewart, 2011).



Opening of Master in Clinical Psychology and Trauma Treatment

At present, 13 students have successfully graduated with a master's degree and many of them are working in private practice as program coordinators and counsellors in the field of trauma and mental health (some with extensive additional training in Eye Movement Desensitization and Reprocessing (EMDR)). Some have started setting up a mobile team serving other provinces. The second cohort of students started its studies in February of 2011 (see: www.masterpsychrupp.webs.com).

As in many other low-income countries, training facilities are still inadequate and professionals are scarce in Cambodia (Saxena et al., 2007). However, a study by Henderson et al. (2005) showed a significant improvement in the confidence of primary care providers in Cambodia in all clusters of medical and psychiatric procedures including e.g. counselling, prescribing medications, psychiatric diagnosis, traditional treatments, and treating trauma victims after extensive training.

Additionally, the donor preference for short-term project-based funding, purely donor driven activities, internal conflicts in mental health leadership, and a lack of a common vision of suitable mental health approaches, hinders the development of sustainable and long-term committed mental health services under Cambodian ownership (Stewart et al., 2010).

Resources in Cambodia

- Up to now 49 trained psychiatrists and 45 psychiatric nurses who work in mental health facilities and/or private practice in Cambodia
- 68 outpatient psychiatric departments (OPD)
- Mental health services available at 9 out of 24 provinces
- 1994 start of undergraduate program at the DP (RUPP) and training of psychiatrists at the University of Health Sciences
- Over 660 students graduated from the undergraduate program in psychology
- 2008: start of Master of Arts in Clinical Psychology and Trauma Treatment at DP, 13 graduated Master of Psychology
- Training facilities still inadequate in most instances
- Scarcity of professionals in Cambodia
- NGOs: leading role in responding to conflicts and disasters by development of new services

Mental health policy in Cambodia

In the field of mental health, professional skills, training, and supervision determine the state of the art and its quality rather than advanced equipment, machines and technology. However, this training is so far not a special focus for the government. According to the Cambodia Health Information System (2007):

"Capacity [in the health sector] is not adequate. The major obstacle to a well performing HMIS [Health Management Information System] is the acute shortage of dedicated and qualified personnel due to low salary and lack of staff motivation."

The majority of civil servants make an estimated \$28 per month, while the Gross Domestic Product in Cambodia is 1,952 USD per capita (UNDP, 2010). The low coverage of mental health

services in the country produces a 0.001 annual rate per capita for utilization of mental health services (Ministry of Health, 2010).

Mental health policy

- Developed countries have a ratio of psychiatric staff per 100000 population that is about 200 times higher than the one in developing countries (Saxena et al., 2007).
- Ranked 124 out of 169 countries in the United Nations Development Program's Human Development Index, Cambodia is a country far from socioeconomic justice (UNDP, 2010).
- The Cambodian budget for mental health is less than 1% of the total annual health budget (Ministry of Health, 2010).

As individuals with mental illnesses are vulnerable to human rights abuse due to disadvantages in cognitive functions, a mental health policy framework must include legislation for protection of their basic rights, especially those in receipt of involuntary treatment (as decreed in Cambodia's National Mental Health Plan 2003-2022) (Saxena et al., 2007). This government-supported national plan envisages positive measures for treatment of mental illnesses and an effective mental health system. However, such plans have so far not been put into action.

Community-based care is even more advantageous for Cambodia as 80% of its population live in rural areas. Although the official mental health policy favours decentralized, community-based care, there are actually no such regular services (Stewart et al., 2010).

Community based mental health care

- Community-based mental health care can be defined broadly, as "any type of care, supervision and rehabilitation of patients with mental illness outside the hospital by health and social workers based in the community" (Saxena, 2007, p. 878)
- From experiences in many countries over the world, a good balance of community-based and hospital-based services has been shown to meet the needs of mentally ill most effectively (Thornicroft & Tansella, 2004) and be more costeffective than long-term treatment in asylums for chronically mentally ill (Saxena et al., 2007)
- 80% of the total of 14.5 M inhabitants in Cambodia live in rural areas and 37% of the population live on less than 1 USD per day (National Institute of Statistics, 2008)

Through the present survey, the DP hopes to raise awareness about the importance of mental health and the relative cost-effectiveness of mental health care (Patel et al., 2007). However, the negative stigma of mental disorders remains strong and hinders the development of adequate and sufficient mental health services. This limits the use of available resources for those in need because they fear social isolation.

Mental health literacy

- In Cambodia many families try to conceal mental illness and hide their ill relatives
- "Educating the public about mental disorders and their treatment is important for improving the empathy of the citizens toward those individuals suffering from mental health problems, reducing discrimination and stigmatization, and improving people's access to general services. It is also important to educate the public on how and when to contact mental health services in order to improve accessibility of services" (MoH, 2003, p. 14)

Cultural explanations for and presentations of mental illnesses in Cambodia

The Khmer concept of individual health is rooted in 4 basic traditional practices rather than a single 'physical' model of health:

Cultural explanations (traditional and moral beliefs) for mental illness

- Buddhist-Hindu beliefs: "Dharma"
- Beliefs in spirits
- Concept of luck (astrology and fortune tellers)
- Somatic/physiological concept
- "Dharma" (correct action) and "Karma" (the balance of good and negative deeds or "fate") are major elements of belief in Cambodia. From this derives a strong spiritual and behavioural code of conduct in Cambodia (Stewart et al., 2010).
- Animistic beliefs include the concept of "unappeased" ancestors as the cause of mental illness. Angry ancestors are placated by ceremonies and offerings placed on the family's shrine.
- 3. The Cambodian belief system incorporates the concept of luck and astrology. Thus, fortune tellers are consulted to assist in preparation for the future. Hinton, Pich, Chhean & Pollack (2005 a, p.55) illustrate this:

"Krauch attributed his frequent khmaoch sångkåt [sleep paralysis] to two facts: 'low bodily energy' (meun mien kamlang dål) and 'low good luck' (rieseuy choh). He explained that if you have 'low good luck' (rieseuy choh), then nothing you attempt to do succeeds; that while in a state of vulnerability (owing to low good luck or weakness), a bad sorcerer (kruu cweng) could send a ghost to attack you."

4. Mental illnesses can result from physical problems (e.g. nausea, palpitations, dizziness, and low energy) and can be treated by easily available medications. "Kyol goeu" (wind overload) is an example of orthostatic irregularity often associated with panic disorder. A Khmer nurse refugee in the USA describes it as: "When wind increases in the body, the person feels aches (especially at the joints), weakness, some anxiousness, and usually a little dizziness. This increase in inner wind may result from eating poorly, from decreased or disturbed sleep, and from thinking too much about one's problems. All of these processes ... weaken the body in a more general way, causing the heart to beat less vigorously than usual, resulting in feeble circulation. Decreased cardiac power results in stagnation and hence further coagulation. Poor circulation also leads to less exiting of wind from the body." (Hinton,

Um & Ba, 2001, p. 409)

The role of culture is a major factor in both the explanation and presentation of mental disorders. The use of cultural narratives and idioms create uniquely Cambodian pathologies (Rechtman, 2000; van de Put & Eisenbruch, 2002). Hinton and others have described a range of key descriptors of Khmer trauma reactions:

- Culturally specific panic attacks, "wind attack" (khyâl) (e.g. orthostatically induced and neck-focused panic) that lie at the centre of the Khmer response to trauma (Hinton, Pich, Marques, Nickerson & Pollack, 2010).
- 2. Sleep Paralysis (SP), in which a demon ghost pushes down (khmaoch sângkât`) upon an individual that is either falling asleep or awakening. The experience causes paralysis, a sensation of a hand on the chest or neck, chest tightness and shortness of breath and is interpreted as a sign of bad luck and imminent danger of dying (Hinton et al., 2005a). SP was clearly associated with trauma and PTSD in Cambodian and other cultures (Hinton et al., 2005a; Ohayon & Shapiro, 2000; Paradis, Friedman & Hatch, 1997).
- 3. "Weak Heart" syndrome (khsaoy beh daung), which involves the belief that "excessive bodily wind" (kyol goeu) causes a breakdown in functioning of the heart with palpitations on slight provocation (e.g., orthostasis, an odour, being startled, exercise-induced palpitations (Hinton, Hinton, Um, Chea & Sak,

2002, Hinton, Hinton, Eng & Choung, 2011; Hinton et al., 2010)). Lambert (1986) has described the "weak heart" as a local manifestation of PTSD, whereas Hinton et al. (2002, p.324) perceive it as corresponding more to the western diagnosis of Panic Disorder.

Treatment seeking behaviour and treatment options for mental illnesses in Cambodia

Until the mid to late 1990s there were no professional counselling services available in Cambodia. The pattern of treatment seeking in Cambodia generally starts with reliance on individual or family coping methods, including:

- · Recreational activities
- Problem solving
- · Support from loved ones
- Drinking only warm water (traditional)
- "Coining" (rubbing a coin on the upper arm or on other parts of the body)
- "Cupping" (suction is created on the skin to mobilizes blood flow in order to promote healing)
- · Herbal remedies
- Alcohol/drug use
- Gambling

In case none of these actions brings relief, Cambodians often seek the assistance of monks, traditional healers (kru khmer), or a medium or fortune teller to alleviate their symptoms through meditative prayer, blessing ceremonies, communication with ancestral spirits (Bertrand, 2005; van de Put & van der Veer, 2005; van de Put & Eisenbruch, 2002). The belief in protection through Buddhist rituals is described by Hinton et al. (2005 a, p. 60):

"Lor visited the local Khmer temple where a Buddhist monk tied a white string around his wrist to prevent the soul from leaving the body and creates a holy, protective barrier."

Western medicines are used when traditional approaches do not alleviate the problem. Neighbours often exchange medication they have been given (Pickwell, 1999); otherwise a pharmacist or

physician is contacted for help. Individuals are traditionally ignorant about the name, ingredient, hoped-for effect and side effects of the prescribed medication.

Because the expectation of many rural Cambodians is to be given at least several medications in order to alleviate their problems, a general habit of over-prescribing drugs has developed all over Cambodia (see: Stewart et al., 2010, McLaughlin & Wickeri, 2012). Additionally many remedies which are prescription-based in the western world are available over the counter without any prescription at all in Cambodia. The low quality of assessment and information and limited availability of appropriate modern medication at many health centres create an increased risk of drug over-use and abuse.

In addition, by the time people finally become aware that more sophisticated treatments are available in Phnom Penh or abroad, most families have already spent a small fortune on local cures. Thus, the vicious cycle of mental illness, discrimination and impoverishment continues.

Inhumane treatments of individuals suffering from schizoaffective or schizophrenic disorders have been reported in many rural areas (Stewart et al., 2010, Ministry of Health, 2003) and observed by staff from the DP. These include putting the affected individual into cages or chaining them, sometimes for months or even years. The study included screening questions for such treatments to better understand how commonly these practices are used and as an indirect indicator for possible psychotic disorders.

Coping and treatment options in Cambodia

- Reliance on family and friends
- Traditional: "coining", "cupping"
- Herbal remedies
- Blessings from and offerings at a pagoda
- Communication with spirits through a medium
- Traditional healers (kru khmer)
- Western medicines with potential danger of harm due to insufficient information
- Many inadequately treated mental illnesses



Self constructed cage in which a client with schizoaffective disorder spent two years because of aggressive and selfharming behavior

Research findings regarding the different issues addressed in this survey

There is little information on the nationwide prevalence of mental illnesses. Most of the studies in this area have so far focused on quite specific aspects or on subpopulations such as refugees. The largest amount of studies has been conducted with Cambodian refugees in the USA. These diverse studies used different measuring scales, addressed different populations, and used different methodology. Their outcomes can only roughly be compared. Additionally cross-cultural challenges of diagnosing mental disorders exist. Many researchers have used the "unverified assumption that DSM-IV disorders have diagnostic validity across cultures" (e.g. de Jong, Komproe & Ommeren, 2003, p. 2128).

An attempt at sorting and clustering the various results according to sub-topics is undertaken in this report. Overall, it can be assumed that Cambodians suffer from significant levels of daily stress, mainly due to low quality of life, poverty, and a much higher percentage of mental disorders (especially depression, anxiety and PTSD) than normative populations (Sonis et al., 2009; de Jong et al., 2003; Dubois et al., 2004). For the general population in Cambodia, these studies estimate prevalence rates ranging from 40 to 53% for anxiety disorders, 11.5 to 80% for

depression, and 7.3 to 86% for PTSD. Comorbidity (i.e. the appearance of several mental illnesses at the same time) is also particularly high (for details see table 1 in the appendix).

1) Violence and mental health

A traumatic experience is defined by the Diagnostic and Statistical Manual IV as an event or events that "involve actual or threatened death or serious injury, or a threat to the physical integrity of the self or others".

Exposure to prolonged or repeated type 2 trauma, common in Cambodia, has a cumulative and far more debilitating impact then single or short traumatic episodes (Field, 2011) and is associated with higher levels of psychopathological conditions (Green et al., 2000; Mollica, McInnes, Pham, Smith Fawzi, Murphy, 1998a; Mollica, McInnes, Poole & Tor,1998b; Killpatrick, Resnick, Saunders & Best, 1989; Follette, Polusny, Bechtle & Naugle, 1996).

Sexual violence against children is prominent in Cambodia (Miles, 2006 al.). According to the World Report on Violence against Children (Pinheiro, 2006), most sexual childhood abuse is committed by family members or their friends in a home setting. "Most children do not report sexual violence experienced at home because they are afraid of what will happen to them and their families (e.g. shame, rejection, abandonment, or disbelief) (Pinheiro, 2006, p 55).

Gang rape seems to be a growing phenomenon in Cambodia. In a qualitative survey conducted by UNICEF (Woods, 2007) and a quantitative Tearfund study (Miles, 2006 b), some young Cambodian men considered it acceptable to gang rape (bauk) girls or women perceived as promiscuous, sexually available, or as prostitutes. A set of questions regarding such experiences was included in the present study.

Violence and mental health

- A traumatic experience is defined as an event or events that "involved actual or threatened death or serious injury, or a threat to the physical integrity of the self or others"
- Exposure to prolonged or repeated, so called type 2- trauma has a cumulative and far more debilitating impact then single or short traumatic episodes
- Most sexual childhood abuse is committed by family members or their friends in a home setting
- Most children do not report sexual violence they experience at home because they are afraid of shame and rejection

2) PTSD

To meet the criteria of a posttraumatic stress disorder (PTSD), according to DSMIV and ICD 10, three categories of symptoms have to be present: persistent re-experiencing, avoidance of stimuli associated with the trauma, and persistent symptoms of increased arousal (see panel 1).

Several studies conducted among the US population show a variety of risk factors associated with PTSD. These include childhood physical abuse (Bremner, Southwick, Damell & Chamey, 1996), trauma in childhood, previous psychiatric history, parents and siblings with mental illness (Bromet, Sonnega & Kessler, 1998), and the severity of trauma exposure (Green, 1990, Kilpatrick & Resnick, 1993). For the PTSD rates in Cambodian populations found in different studies, please see table 1 in the appendix.

Additionally, the ongoing challenges in coping with psychological consequences stemming from the KR regime, such as living with survivor's guilt and experiencing feelings of revenge have been examined and reported in various studies (Pham, Vinck, Balthazard, Hean, 2011; Kirchenbauer, Burchert, Taing, Bockers, Knaevelrud, 2010; Rechtmann, 2006; Field & Chhim, 2008).

Panel 1 DSMIV-TR criteria for PTSD

- A. The person has been exposed to a traumatic event in which the following were present:
 - 1. The person experienced, witnessed, or was confronted with an event or events that involved actual or threatened death, or serious injury, or a threat to the physical integrity of self or others
 - 2. The person's response involved intense fear, helplessness, or horror.
- B. Persistent re-experiencing in one of the following ways:
 - Recurrent and intrusive distressing recollections of the event, including images, thoughts, or perceptions.
 - 2. Recurrent distressing dreams of the event. In children, there may be frightening dreams without recognizable content.
 - 3. Acting or feeling as if the traumatic event were recurring (includes a sense of reliving the experience, illusions, hallucinations, and dissociative flashback episodes, including those that occur when awakening or intoxicated). In young children, trauma-specific re-enactment may occur.
 - 4. Intense psychological distress at exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event.
 - 5. Physiological reactivity on exposure to internal or external cues that symbolize or resemble an aspect of the traumatic event.
- C. Persistent avoidance of stimuli associated with the trauma and numbing of general responsiveness (not present before the trauma), as indicated by three or more of the following:
 - 1. Efforts to avoid thoughts, feelings, or conversations associated with the trauma
 - 2. Efforts to avoid activities, places, or people that arouse recollections of the trauma
 - 3. Inability to recall an important aspect of the trauma
 - 4. Markedly diminished interest or participation in significant activities
 - 5. Feeling of detachment or estrangement from others
 - 6. Restricted range of affect (e.g. unable to have loving feelings)
 - 7. Sense of a foreshortened future (e.g. does not expect to have a career, marriage, children, or a normal lifespan)
- D. Persistent symptoms of increased arousal (not present before the trauma), as indicated by two or more of the following:
 - 1. Difficulty falling or staying asleep
 - 2. Irritability or outbursts of anger
 - 3. Difficulty concentrating
 - 4. Hyper-vigilance
 - 5. Exaggerated startle response
- E. Duration of the disturbance (symptoms in Criteria B, C, and D) is more than 1 month.
- F. The disturbance causes clinically significant distress or impairment in social, occupational, or other important areas of functioning.

Cambodians seem to experience hyper-arousal mainly in the form of dizziness and blurred vision whereas blocked traumatic energy leads to chronic muscle tensions causing headaches, backaches, and stiff necks. In addition, many KR survivors experienced physical trauma and still suffer from vivid body memories or chronic physical lesions.

3) Cultural validity of the PTSD construct

In the world's cross-cultural trauma literature, the western concept of PTSD is disputed as ignoring somatic and dissociative symptoms and focusing too much on avoidance (Berthold & Gray, 2011C). Research in Asian and African nations have created new tools and are advocating for the inclusion of more cross-cultural aspects in the new version of the DSM and found considerable evidence of the cultural structural and content

validity of PTSD (Hinton & Luis Fernández, 2010).

It was also found that "many of the 17 PTSD items listed in the DSM IV manual, such as nightmares, startle reflexes, and vivid unwanted recall of trauma events are a core part of the universal response to trauma. [However] other DSMIV PTSD items like amnesias or numbing seem to be much less salient aspects of the trauma response in non-Western cultures." (Hinton et al., 2011, p. 51). In a study on sleep paralysis, which is not mentioned in Western concepts of PTSD, Hinton et al. (2005 a) found a much higher prevalence of SP in clients with PTSD than without probably due to the high arousal level. This shows the importance of using culturally adapted versions of assessment tools and instruments especially developed for certain cultures in research.

4) Anxiety and depression

"Gender determines the differential power and control men and women have over the socioeconomic determinants of their mental health and lives, their social position, status and treatment in society, and their susceptibility and exposure to specific mental health risks." (WHO, 2011 c). Unipolar depression may become the leading cause of the global disability burden by 2020 (WHO, 2011 c). The Global Burden of Diseases (GBD) 2000 estimates a point prevalence for unipolar depression of 1.9% for men and 3.2% for women and a 12 months prevalence of 5.8% for men and 9.5% for women (WHO, 2001).

In a study by Jegannathan & Gunnar (2011) with 320 high school students around Takhmau aged 15 to 18, young women scored higher on anxious/depression, withdrawing/depression, somatic complaints, social problems, and internalizing syndrome whereas young men scored significantly higher on rule-breaking behaviour than young women.

Governmental estimates of percentages of the most common mental disorders treated at governmental health centres in Cambodia are: anxiety disorders (39%), depression (29%), psychosis (18.5%), epilepsy and organic problems (5.2%), substance abuse (3.0%), and reactions to severe stress and adjustment disorders (1.9%) (Stewart et al., 2010). For other study results in Cambodia please see table 1 in the appendix.

Anxiety and depression

- "Gender differences occur particularly in the rates of common mental disorders depression, anxiety and somatic complaints. These disorders affect approximately 1 in 3 people in the community and constitute a serious public health problem." (WHO, 2011c)
- Worldwide depression is the most common mental health problem in women

5) Tendency to suicide

"In the last 45 years, suicide rates have increased by 60% worldwide" WHO (2011a). Suicide is among the three leading causes of death among those aged 15 - 44 years-olds in some countries. Although suicide rates have traditionally been highest among the male elderly, rates among young people have increased to such an extent that they are now the group at highest risk in a third of countries, in both developed and developing countries. Mental disorders (particularly depression and alcohol use) are a major risk factor for suicide in Europe and North America; however, in Asian countries, impulsiveness plays an important role. Suicide is complex with psychological, social, biological, cultural, and environmental factors involved." (WHO, 2011a).

"As of May 2003 the highest rates for accomplished suicides were found in eastern European countries such as Lithuania, Hungary, Russia, Slovenia, Kazakhstan, and Estonia. High rates have also been recorded for Japan and Sri Lanka." (WHO, 2011b).

In the US, suicide is the eleventh leading cause of death for all US Americans, and the third leading cause of death for young people 15 to 24 years old. A study by Weissman et al. (1999) confirms these findings and describes suicide attempts as relatively consistent across nine countries from North and South America, Europe, Near East, Asia, and New Zealand. They also found that attempts in most countries were associated with being currently divorced or separated as compared to currently married.

There exist many examples of secretly desperate people driven to attempt suicide in Cambodia. Newspapers report many attempted and committed suicides (Cambodia daily, 2011). Severe trauma, torture, and refugee experiences were associated with higher suicide rates and 40% of refugees from different countries with PTSD attempted suicide (Ferrada-Noli, Asberg, Ormstad, Lundin & Sundbom (1998)). Suicides and suicide attempts have therefore received special attention in the present study.

Facts on suicide

- Every year, almost one million people die from suicide (WHO, 2011a)
- "Global" mortality rate of 16 per 100,000, or one death every 40 seconds from suicide (WHO, 2011a)
- Suicide is the second leading cause of death in the 10-24 years age group (WHO, 2011a)
- Suicide attempts are up to 20 times more frequent than completed suicide (WHO, 2011a)
- In the US males are four times more likely to die from suicide than females. However females attempt two to three times more often to commit suicide (Centers for Disease Control and Prevention (CDC), 2009)
- Compelling evidence indicating that adequate prevention and treatment of depression and alcohol and substance abuse can reduce suicide rates, as well as follow-up contact with those who have attempted suicide (WHO, 2011a)

6) Second generation trauma

At present, over 60% of Cambodia's population was born after the KR regime (Munyas, 2008). "The loss of family ties, rights, dignity, and honour has been in the heart of the Cambodian population for a full generation. Due to this troubled history, many parents did not have the experience of positive parenthood. Instead, war and public violence was a grim reality faced on a daily basis. As a result, a whole generation of Cambodians is missing parenting skills and models for the peaceful resolution of conflicts" (Woods, 2008, p.10). In Cambodia, typically a parent-child role reversal (or overprotective behaviour) has been found deriving from the large prevalence of traumatic experiences (Field, 2011).

This may be a critical factor explaining the high level of violence experienced by the second generation of Cambodians after the KR era. According to Field (2011, p. 77):

- 47% experienced physical punishment by parents
- 14% were sexually abused
- 22% witnessed the rape of another child by an adult
- 48% had knowledge of a child having been sold

These findings also suggest that traumatized parents failed to protect their children and probably caused physical and emotional neglect. Such second generation relational trauma is only indirectly visible and thus often ignored.

A study by de Jong (2001) revealed a significant predictive value of domestic stress for PTSD in a Cambodian sample, which was not present in the three other countries examined (Algeria, Ethiopia, and Gaza).

As a result, the present study included screening questions about child and adolescent mental health and problematic behaviours.

Second generation

- Today over 60% of Cambodia's population was born after the KR regime
- Due to the nation's troubled history many of today's parents did not have the experience or example of positive parenthood
- High level of violence experienced by the second generation of Cambodians after the KR era
- Findings suggest failed protection by traumatized parents and probable physical and emotional neglect

7) Addictive disorders

"The lifetime prevalence rate for alcohol dependence is more than twice as high in men than women. In developed countries, approximately 1 in 5 men and 1 in 12 women develop alcohol dependence during their lives." (WHO, 2011c). According to the GBD (2000) the point prevalence for harmful use of or alcohol dependency is 1.7% overall, 2.8% for men, and 0.5% for women (WHO, 2001). Prevalence rates in a representative sample from the USA show 17.8% lifetime alcohol abuse, 12.5% lifetime alcohol dependence, 4.7% 12 month alcohol abuse and 3.8% 12

month dependency (Hasin, Stinson, Ogburn & Grant, 2007).

Addictive disorders often develop as a sort of coping strategy for dealing with overwhelming emotions, often abundant and horrific fears, social isolation, stigmatization and loss of capacities, or low self-esteem.

Many people with psychological problems resort to self-medication with drugs, alcohol, or over the counter medication that can be very harmful. Others take up gambling. A meta-analysis by Stewart (1996) shows a significantly elevated risk for the development of alcohol-related problems for individuals suffering from PTSD when compared with both individuals exposed to trauma without PTSD and those who have never been exposed to a traumatic event. This is suspected to be due to self-medication with alcohol for PTSD symptoms although, in the long term, alcohol seems to increase symptoms of anxiety and PTSD rather than being helpful.

Addictive disorders may create a bad model for youth and influence them to engage in such behaviour. This is particularly alarming with respect to sniffing glue and amphetamine abuse in Cambodia (Crookes, 2011; UN, 2010). Therefore a set of questions regarding addictive behaviours was added to the questionnaire of this study.

Addictive disorders

- The lifetime prevalence rate for alcohol dependence is more than twice as high in men as in women
- In developed countries, approximately 1 in 5 men and 1 in 12 women develop alcohol dependence during their lives
- According to the GBD 2000 the point prevalence for harmful use of or alcohol dependency is 1,7% overall, 2,8% for men and 0,5 % for women
- Addictive disorders as a coping strategy for dealing with overwhelming emotions, fears, social isolation, stigmatization, loss of capacities and low self-esteem

8) Domestic violence

The Law on Prevention of Domestic Violence and Protection of Victims (Kingdom of Cambodia, 2005) states in Article 2 that, "Domestic violence as referred to the violence that happens and could happen towards: husband or wife, dependent children, persons living under the roof of the house and who are dependents of the households."

Domestic violence can be divided into physical, sexual, emotional and financial violence. The Cambodian League for the Promotion and Defence of Human Rights (LICADHO) notes "a strong relationship between violent conflict in society and violence in the home. [For example,] after three decades of war, violence is an accepted end to conflict in most of Cambodia." (Pact Cambodia, 2004).


Communication skills that might prevent misunderstandings and domestic stress seem to have a low cultural value and are widely underdeveloped in Cambodia. Opinions therefore are seldom shared and points of view are not explained.

The incidence of domestic violence remains high, affecting 20 to 25% of women (MoWA, 2008). The analysis of the Cambodian Demographic and Health Survey 2005 showed that women earning more are at higher risk for mental violence whereas less educated women were associated with greater physical domestic violence. (For results on this see: full report)

9) Schizophrenia

"There are no marked gender differences in the rates of severe mental disorders like schizophrenia and bipolar disorder that affect less than 2% of the population" (WHO, 2011c). There are no studies available on schizophrenia or psychotic disorders in Cambodia so far. The world health report 2001 mentions a point prevalence of 0.4% according to the GBD 2000 (WHO, 2001).

As mentioned before, this disorder has severe consequences and often affects the dignity and human rights issues of individuals. Therefore indirect indicators for this disorder have been included in our screening part of the questionnaire.



A nationwide mental health survey was conducted among 2,690 adults from nine provinces and Phnom Penh in Cambodia (July - August, 2011). The study was reviewed and approved by the National Ethics Committee for Health Research.

I Sample

The sampling technique and the sample size were decided upon based on a Cambodian study on violence against women conducted by the Ministry of Women's Affairs in 2009 (GTZ, 2009). This research targeted a nationally representative sample of Cambodian adults (21 years of age and above) living within Cambodia by using a multi-stage cluster sampling technique. ity proportional to size (PPS).

The specifics of the multi-stage cluster sampling for this research included five stages:


- 1: Cambodia comprises 23 provinces that can be grouped into four geographic regions (i.e. plain, coastal, plateau/mountain, and Tonle Sap). Two provinces per region were selected nine provinces in total (incl. three provinces for Tonle Sap). The capital of Phnom Penh was also integrated. Thus, the nine selected provinces & Phnom Penh comprised Koh Kong, Preah Sihanouk, Kampong Speu, Kratie, Prey Veng, Takeo, Phnom Penh, Banteay Meanchey, Kampong Chhnang, and Pailin.
- 2: Each province was divided into administrative districts. Of these 50 districts were randomly selected using the PPS method (please note: throughout the study, the adult population was determined by the '2008 General Census').
- 3: Based on the PPS method, 100 communes were selected within the districts. The communes in the selected districts were further stratified into urban and rural (14% were urban communes).
- 4: 270 villages (2 to 3 randomly selected from each stratum) were identified from the selected communes.
- 5: Ten households were selected from each targeted village based on a fixed interval according to the total number of families in the village and a 10:1 sampling ratio. Once a household was identified, the Kish Grid technique was used to select adult respondents in the household for interview. Thus, a total of 2,700 respondents were targeted.

Il Instruments, scales and survey development

The interview questionnaire was divided into 8 parts:

Structure of the survey

- Biographical Background
- Screening Questionnaire
- Domestic violence (National Institute of Public Health, 2005)
- Traumatic events during and after the KR (HTQ) (Mollica et al., 1996), and other stressful events
- Harvard Trauma Questionnaire-Symptoms Part (HTQ), (Mollica et al., 1996)
- The Hopkins Symptom Checklist 25 (HSCL-25) (Mollica et al., 1987)
- Cambodian Addendum of Culturally Sensitive Items, now: Cambodian Somatic Symptom and Syndrome Inventory (C-SSI) (Hinton et al. 2012)
- Questions related to needs and services



2-3 villages randomly selected from each stratum (PPS)

 \rightarrow 270 villages

10 households within each targeted village by PPS

→ 2700 households

Kish Grid technique to select adult respondents in households

In order to deal with the high illiteracy rate, all questions included in the questionnaire were conducted as part of the interview.

The assessment of the biographical background, the screening questionnaire, as well as the questions concerning the availability of psychological services within the target area were developed by a multi-professional team in Cambodia, a Cambodian sociologist (PhD from Mississippi state University) and a psychiatrist (GIZ German Peace Service advisor).

Additional in-depth questions explored reasons

Background information

- Age
- Sex
- Marital status
- · Position in the household
- Years of education
- · Family size
- Number of respondent's children working or attending school
- · Family income and spending





Screening questions Regarding children:

- Antisocial behaviour and contact with police/ court/reeductation camp
- · Aggressive behaviour
- School problems
- Other problems

Regarding respondents and family:

- Indicators for psychotic or schizoaffective disorders (including being held in a cage/ being chained)
- Aggressive behavior
- Suicide and suicide attempts
- Violence experienced
- Addictive disorders

for suicidal behaviour, substance abuse, and other possible addictive behaviours. This screening part of the interview (developed in English

and Khmer) used typical Khmer expressions for some phenomena. Items developed in English were extensively discussed for an accurate translation by field supervisors.

Questions on 'domestic violence' were selected from the Cambodian Demographic Health Survey (National Institute of Public Health, 2005) assessing violent behaviour conducted or experienced in a domestic context as well as the interviewee's attitude towards domestic violence.

The HTQ, HSCL-25, and the CSSI are commonly used instruments in research conducted in Cambodia and humanitarian contexts (Dubois et al., 2004). "The HTQ and HSCL-25 were initially developed to assist clinicians in assessing the mental health of refugee patients in specialist refugee mental health services and in primary care settings" (Mollica, McDonald, Massagli & Silove, 2004, p. 4). They have been translated, backtranslated and validated extensively (Sonis, 2009) and address only the respondent.

The first part of the HTQ assesses traumatic events during the Khmer Rouge regime such as torture, witnessing murder, and lack of food and water. The original list was adapted to address Cambodian experiences. The number of different traumas out of 20 was summed to create an index of trauma experienced during the KR era.

Two lists of events were added to the HTQ. The first list assesses traumatic events that occurred after the Civil War, such as severe accidents, natural disasters or rape. In this part respondents are asked whether they experienced or witnessed the mentioned traumatic events. The number of different traumas, out of 12 was summed to create an index of trauma experienced after the KR era.

The second list included stressful events happening within the families such as severe illnesses, death of a close relative or severe financial problems. The number of different stressful experiences, out of 13 was summed to create an index of stressful life experiences.

The second part of the HTQ includes 16 items as criteria for PTSD that were derived from the Diagnostic and Statistical Manual of Mental Disorders

(Third Edition, Text Revision) (DSMIII-R). Respondents were asked to rate the intensity of the symptoms on a Likert scale from 1 ("not at all") to 4 ("extremely"). The threshold-score for PTSD was evaluated as ≥2.50. Validity and Reliability of the HTQ are sufficiently high. The term 'probable PTSD' was used because the HTQ, despite good measurement properties, is not a diagnostic interview. In the current study, the HTQ-Symptoms had excellent internal reliability (Cronbach = 0.91).

HTQ-Symptoms

- 16 items that derive from the DSMIII-R criteria for PTSD
- Likert scale for the intensity of the symptoms from 1 ("not at all") to 4 ("extremely") true
- The threshold-score for PTSD evaluated as ≥2.50 for Southeast Asian patients
- Validity and reliability of the HTQ sufficiently high (internal reliability in this study: Cronbach's alpha = .91)

The HSCL-25 consists of 25 items measuring anxiety, depression, and somatic symptoms. Respondents were asked to rate the intensity of the symptoms on a Likert scale from 1 ("not at all") to 4 ("extremely true"). Subjects with a score >1.75 on the depression scale were classified as probably depressed whereas subjects with a score >1.75 on the anxiety scale were classified as probably suffering from anxiety. Quality criteria are sufficiently high: Sensitivity and specificity of the Indochinese version of the HSCL-25 are around 0.88. 'Retest reliability' for the total score is 0.89 and 0.82 for the depression score. Interrater reliability was estimated as 0.98. In the current study, the HSCL-25 had a very good internal reliability of Cronbach = 0.86 for anxiety and 0.86 for depression. Again we use the term probable anxiety or depression because the HSCL-25 is not a diagnostic interview.

The Hopkins Symptom Check-List-25

- 25 items (10 items for anxiety, 13 items for depression, 2 items for somatic symptoms)
- Depression and anxiety scores
- Likert scale for the intensity of the symptoms from 1 ("not at all true ") to 4 ("extremely true")
- Threshold scores in a Cambodian Context: >1.75 for depression, >1.75 for anxiety
- Sensitivity and specificity .88
- Retest-reliability for the total score is .89 and .82 for the depression score
- Inter-rater reliability was estimated .98
- Internal reliability in this study .86 for anxiety and depression

The Cambodian Somatic Symptom and Syndrome Inventory (C-SSI) is a tool to assess typical (often trauma related) symptoms in a Cambodian population. The C-SSI was previously called the "Cambodian Addendum of Culturally Sensitive Items".

The C-SSI consists of two parts. The first part measures culturally predominant somatic symptoms that correspond partly to anxiety symptoms and are missing in the Western PTSD description. The second part measures key cultural syndromes and constructs often related to typical causal attributions (e.g. evil spirits, ghosts, and sleep paralysis caused by *khmaoch sângkât*). Respondents were asked to rate the intensity of the symptoms on a Likert scale from 1 ("not at all") to 4 ("extremely true") referring to the previous month.

The Cambodian Addendum of Culturally Sensitive Items has been used frequently with Cambodian refugees (Hinton et al., 2005). In the current present study it had excellent internal reliability (Cronbach = .89).

The Cambodian Somatic Symptom and Syndrome Inventory (C-SSI)

- Typical (possibly trauma related) symptoms in a Cambodian population
- 2 parts:
 - Culturally predominant somatic symptoms that are missing in the PTSD concept
 - Key cultural syndromes and constructs often related to typical causal attributions
- Likert scale for the intensity of the symptoms from 1 ("not at all true") to 4 ("extremely true")
- Maximal total score of 4.29
- Easily understood by interviewees

Questions on 'mental health services available' were developed by the research team and covered the accessibility and utilization of mental health services in the area of the respondent as well as general knowledge about psychology.

Needs and services

- Accessibility of mental health services in the area of the interviewee
- Who is contacted by the interviewees in case psychological problems occur





General knowledge of psychology

III Data collection procedures

Face to face surveys were administered by students from the Department of Psychology at the Royal University of Phnom Penh (13 males and 13 females). Five field supervisors (all with master's degrees in psychology) implemented strict quality control, including detailed review of completed interviews and random observation of interviews in progress during the survey. All members of the data collection team received 19 hours of training in sampling, interviewing skills, and basic trauma response skills. The training was provided to the DP by a GIZ advisor (German psychiatrist and expert in traumatology) and a statistician (member of the National Committee for Population and Development with a PhD in Sociology).

Data collection teams consisted of five interviewers and one supervisor. An average of 4 interviews was conducted per day. Each interviewer conducted approximately 110 interviews (each interview lastinged between 40 to 100 minutes). Distances between villages were often very far. Only 18 interviews were not completed. Although strict confidentiality was intended, approximately 50% of the interviews were conducted with other household members present (data collectors evaluation report). Interviewers were asked to fill out an evaluation questionnaire concerning their experiences and possible problems occurring during data collection.

All respondents were informed about the subject and intent of the interview. Due to the high rate of illiteracy and fear of signing forms, respondents gave oral rather than written consent to participate in the study. No incentives were offered to respondents. If interviewees were assessed as needing psychological or psychiatric counselling, such therapy was provided treatment was planned for offered via the Mekong project. The project provides free treatment for traumatized individuals via HAP (Human Assistance Program - Germany).

Data collection

- Interviewers: 25 undergraduate and graduate psychology students
- 5 field supervisors (master in psychology)
- Data collection teams were trained in intervierwing skills and basic trauma response skills (2 days)
- Pilot testing
- Feedback-questionnaire and debriefing meeting after the data collection
- Data collection teams : 5 interviewers & one supervisor
- An average of 4 interviews per day per data collector
- Duration of interviews: 40 to 100 minutes
- Oral consent to participate in the study
- No incentives
- 18 interviews uncompleted

IV Data entry

Double data entry was performed by 7 experienced psychology students between late August and late October. All were trained by the project assistant. The data was checked for data entry errors using Epilnfo and errors were corrected accordingly. Further checks for inconsistencies and corrections were conducted by the team's statistician, the German advisor, and the team. Some misinterpretations were detected due to translational issues (e.g. the English word "children" has two different translations and meanings referring to one's own and other children in Khmer).



V Data analysis

Data analysis was conducted using SPSS 16 and sometimes used a multi-level model when splitting data concerning rural and urban areas, sex, and age. Grouping into rural and urban areas was done to acknowledge the possibility of different experiences of the respondents during the KR regime as well as natural disasters and/or traffic accidents.

Sex (as a grouping variable) was included as many studies have found significant differences in symptoms and syndromes of mental health problems in men and women (de Jong, 2003).

Age was used as a grouping variable as the authors expected to find different outcomes for people who have directly experienced the KR and the second generation who might be indirectly affected by traumatized parents and/or other community elders. Therefor interviewees were categorized as those over and those under 36-years-old. Respondents under the age of 36 had to be at least 3 years old at the end of the KR regime.

Background information, mental health services, and the screening questionnaire were analyzed descriptively. For the screening questionnaire children's problems and problematic behaviours were estimated per household with respondents having children (N = 1848) and per respondents' children (N = 4002). It was also differentiated between members of the household (excluding the respondent) and problems of the respondents themselves. This was done as follows:

- Mental disorders affecting the respondent and family members
- Occurrences of traumatic events (such as rape, gang rape, trafficking, child abuse) per household
- Suicide attempts of respondents and their family members
- · Committed suicides per household
- Addiction problems such as alcohol, drugs and gambling

Alcohol consumption was considered to be a probable addiction problem if the respondent reported to drink more than half a litre or two cans of alcohol per day. Alcohol consumption was

considered to be a probable abuse problem if the respondent reported to drink more than a liter or 5 cans of alcohol per week and to have financial problems because of this.

Further analysis using cross-tables was conducted and checked for significance for such variables as alcohol problems, suicide attempts, and the event of being chained or put in a cage. Each variable addressed, on one side, the respondents themselves and, on the other side, demographic data (i.e. age, education, economic, and marital status).

In the trauma-related part of the questionnaire, events witnessed and experienced were combined into one category in order to estimate the total amount of traumatic events experienced by the respondent. The total amount of traumatic events experienced was summed up separately for trauma related to the CW and the ones after the CW. The total of stressful life experiences ("current events") was calculated accordingly. Additional summative scores were calculated for traumatic experiences during and after the CW and the total amount of traumatic and stressful experiences.

For the other symptom scales, summative scores with means (M), standard deviations (SD), and percentages of probable diagnosis - according to the above mentioned threshold scores with confidence intervals (95% CI) - were calculated for each of the total female and male (sub) populations. Internal reliability of all scales was analyzed.

Data was checked for distribution with the Kolmogorow Smirnow test and mostly did not follow normal distribution. Differences in categorical variables across population groups (e.g. sex, education, marital status) were therefore assessed by nominal Phi. Spearman correlations and calculated for traumatic experiences, symptom scales, and continuous demographic variables (age, savings per capita, and years of education). Differences in means of the various symptom scales across population groups were assessed using the Mann Whitney U Test for

dichotomous variables and the Kruskal Wallis H-Test for categorical variables.

Questions about domestic violence were analyzed descriptively and separately for violence

Data entry and analysis

- Double data entry
- Data check for data entry errors with Epilnfo
- SPSS 16
- Data was split up into rural and urban areas, sex and age (under and over 36)
- Data was analyzed mainly descriptively and with Spearman r correlations

committed and experienced by male, female, and all respondents (see full version of report).



Note: All numbers, reported in results, are valid percentages (excluding missing values) The full version of this report will be soon available on: www.masterpsych-rupp.webs.com

I Biographical background - respondent profiles (gender, age, position)

Of the respondents:

- 67.5% female, 32.5% male
- 44.1% = head of household
- 40.0% = husband/wife of head of household
- 10.9% = children of the head of household
- 4.9% = other relatives




Fig. 1: Household Position of Respondents in % (N = 2,671)

Marital status






Fig. 2: Marital status of respondents in % (N= 2,677)

Marital Status	Urban	Rural	Total
Married	602	1,482	2,084
Single	105	94	199
Divorced	36	15	51
Widowed	115	227	342
Total	858	1,818	2,676

Table 1: Marital status and number of respondents by region of origin

Education

Years of education:

Average: 4.96

Minimum: 0

Maximum: 20

• SD: 3.90

Occupation





No education

Primary school

High school

Secondary school

University or collede

Main Employment

- 49.6% Farmers · 4.7% Civil servants
- 18.0% Vendors 0.3% NGO staff
- 12.8% Housewives 0.4% Company staff
- 7.9% Labourers 1.6% Students
- 4.8% Unemployed



2.6

18.1

47.6

10

21.7




Fig. 3: Educational status of respondents in % (N=2,685)

Fig. 4: Main occupation of respondents in % (N= 2,680)

Unemployed

Family members

- Average of 5.31 members per family (Min 1, Max 22, SD 2.08, N = 2688) with 1.49 children per total respondents
- Of all respondents, 69% had children (i.e. 1,848 out of 2,678 respondents)
- Thus 2.17 children per all respondents with children (Min 1, Max 9) of these children, 69% at tended school and 30.4% were employed

Economic situation

Income, spending and savings:

On average, 2.17 people per household earned an income (Min 0, Max 9, SD 1.06, N = 2,671)

Average monthly income:

- For all households: 153.06 USD (SD=137.11)
- For 48.2% of households: less than 50 USD
- For 1.2% of households: greater than 800 USD
- For 0.6% of households: no income was earned

Average monthly spending:

• For all households: 140.56 USD (SD=114.57)

Average monthly savings:

- For all households: 2.53 USD (SD=18.82) (N = 2,681)
- For 87.4% of households: no savings subsistence living only

Average monthly debt incurred over income earned:

For 6.5% of households: greater than 150

Department of Psychology, Royal University of Phnom Penh

The following table shows the economic situation of respondents by location.

Area	Income per household	Spending per household	Savings per capita
Rural	124.15	115.05	2.04
Urban	214.59	194.87	3.60
Total	153.06	140.56	2.54

Table 2: Financial situation of the respondents relative to demographics and average monthly income

Respondents in urban areas earned and spent more, and could save nearly twice as much as rural residents.

Ownership of property and landholdings

Approximately 91.5% of respondents owned land (N = 2,683).




Fig. 5: Amount of land owned by the respondents in % (N = 2,441)



Real assets of the respondent's families:

- 95.8% owned a house (N = 2,686)
- 64.5% owned a motorcycle (N = 2,686)
- 7.3% owned a car or truck (N = 2,681)

Type of dwelling	Percent of type owned
Wood house with tin roof	53.8%
Wood house with tiled roof	32.4%
Thatched house	7.9%
Brick house	5.9%

Table 3: % of resp. with different types of houses (N = 2,555)

II Screening questions

Children problems



The following results concerning children in the families are calculated referring to the selection of respondents having own children, which were 1848 respondents out of the 2690, if not mentioned otherwise.

1) School problems

The following graph shows the percentage of households with children having problems at school.




Fig.6: Percentage of households with respondents' and other children having problems at school (N = 1653/1667 (other/own))

- An average of .09 (SD = .36) households had school problems regarding respondents' children and .02 (SD = .178) regarding other children living in the household
- In total 8.2% of the households had children with school problems

The following table shows the percentage children having school problems in given time frames.

	School problems	School problems
	% within previous week	% within previous month
Respondents' children	25.8	63.7
Other children	43.3	70.0

Table 4: Cumulative percentages of respondents who experienced school problems in certain time frames (N = 30/124 (other/own)

Around two thirds of these children experienced school problems within the previous month

2) Aggressive behavior

The following graph shows the percentage of households with aggressive children.




Fig 7: Percentage of households with respondents' and other children showing aggressive behavior (N = 1,666/1,787 (other/own))

- 9.9% of respondents reported about 1 to 6 of their children showing aggressive behavior (M = .14,
 SD = .474), out of which 78.4% exhibited the behavior during the previous month
- In total 11.5 % of households were dealing with aggressive children in their family

The following table shows the percentage children with aggressive behavior in given time frames.

	Aggressive behavior	Aggressive behavior		
	% within previous week	% within previous month		
Own children	45.5	78.4		
Other children	56.2	79.2		

Table 5: Period in which respondents and other children in the households exhibited aggressive behavior (N = 48/146 (other/own))

Nearly 80% of children's aggressive behavior occurred within the previous month

3) Other problems

The following graph shows the percentage of households with children suffering from other problems.




Fig. 8: Percentage of households with respondents' and other children suffering from other problems (N = 1671/1774 (other/own))

- In total 4.9% of the households had other problems with the respondents children (M = .06, SD = .332), out of which 58.6% occurred during the previous month
- 1% reported about other problems with other children from their families

4) Detention of children

The following graphs present the percentage of households with respondents' own or other children who were brought to a reeducation camp or to police/ court or prison.




Fig.9: Percentage of households with respondents' or other children being brought to reeducation camps (N = 1672/1778 (other/own))

- 3 respondents (0.2%) reported about each one own child that had been brought to a reeducation camp out of which 33.3% within the previous year
- 6 respondents reported about one other and one about 6 other children in the household that had been brought to a reeducation camp, 57.1% within the previous year
- In the whole sample of 2690 households (N= 2365) 22 cases of other children in 9 of the respondents' households had been brought to a reeducation camp. Out of these 33.3% within the previous month and 55.6% within the previous year




Fig.10: Percentage of households with respondents' or other children which had contact with the police, court or went to prison (N = 1659/1782 (other/own))

- In 10 households (0.6%) each one own child of the respondent had been brought to police, court or prison, 60% of these within the previous year
- In 5 households one to three other children had been brought to police, court or prison, 40% within the previous year

5) Percentage of respondents' children suffering from different problems

The following graph shows the percentage of the respondents' children suffering from the above problems calculated in relation to a total number of 4002 children of respondents.




Fig.11: Percentage of respondents' children suffering from various problems

• 5.9% of the respondents' children were exhibiting aggressive behavior, 3.8% problems at school and 2.9% other problems

Traumatic events experienced

- In 0.4% of the households 1 family member had experienced rape, 45.5% within the previous year (N = 2685)
- In 0.7% of the households one to four members of the family had experienced trafficking, 61.1% within the previous year (N = 2684)
- 8.8% of the respondents had experienced child abuse, described as physical or sexual violence or severe neglect, 10.6% of male respondents (M = .01, SD = .03, N = 871) versus 8.0% of female respondents (M = .08, SD = .27, N = 1807)
- In 4.0% of the households one to six other family members had experienced child abuse (N = 2666
- In 5.9% of the households one to five members had experienced another crime (N = 2681), out of which 24.2% within the previous year

Mental disorders of family members

- 93 respondents (3.5%) reported having one to two family member affected with "craziness" and "talking nonsense" (N = 2681)
- An average of .04 members per family were affected (SD = .20)
- 25.8% of these where acute within the previous week, 38.7% within the previous year they were ill for 2 months to 70 years with an average of 16.91 years (SD 14.23)
- Additionally four cases of family members from 0,1% of respondents were reported having been chained or put in a cage because of a mental problem, two of them within the previous month
- 8 respondents (0.3%) reported having been chained or put in a cage because of a mental problem themselves, 0.6% of male respondents (M = .01, SD = .08, N = 870) versus 0.2% of female respondents (M = .00, SD = .04, N = 1811)

- One reported this happened the week before the interview took place
- Together with severe mental disorders of family members this makes 0.74% of the household members affected and a point prevalence of about 0.2%

Irritability/ aggressiveness

- An average of .56 (SD = .496) respondents (56.1%) reported being irritable or angry (N = 2653), 60.4% of female respondents (M = .60, SD = .49, N = 1790) versus 47.3% of male respondents (M = .47, SD = .50, N = 859); 60.4% out of these within the previous month
- 24% of the respondents reported having one to seven family members behaving irritably or angry; 64.9% of these within the previous month

Suicidal tendency

1) Suicide attempts

- 115 (4.3%) respondents reported that they had previously attempted to commit suicide (M= .04, SD = .203, N = 2684)
- 15 were male (1.7% of male respondents, M = .02, SD = .13, N = 871), 100 female (5.5% of female respondents, M = .06, SD = .23, N = 1809) with differences on a highly significant level
- 46 were 36 years old or younger (5% of younger population), 69 older than 36 (3.9% of the older population) (not significant)
- Suicide attempts correlated negatively on a significant level (p<.05) with savings per capita, meaning that respondents with fewer savings committed more suicide attempts
- In 22 cases (0.8%) the respondents reported a family member had attempted to commit suicide (N = 2668)

The following graph shows the percentages of attempted suicides per time frame.




Fig.12: Percentage of attempted suicides within a given time frame, cumulated percentages (N = 115/22 (respondents/ family members))

- 13 out of the 115 respondents (11.3%) had attempted to commit suicide during the week previous to the interview
- 26 (22.6%) during the month previous to the interview
- In total 44.3% of the 115 respondents and 50% of the 22 family members had attempted to commit suicide during the previous year

The following table shows mean scores of different symptom scales for respondents that had or hadn't committed suicide attempts. The differences between the two groups are highly significant (Mann-Whitney-U-Test, p < .001) for all scales.

Suicide attempts	HTQ-Symptoms		HSCL-25 depression	CSSI
No	1.40	1.49	1.42	1.74
Yes	1.91	1.98	1.94	2.22

Table 6: Mean symptom scores related to suicide attempts

 Respondents that had attempted to commit suicide suffer from much higher levels of psychological symptoms

2) Committed suicides

- In 22 households (0.8%) a family member of the respondent had committed suicide (M .01, SD = .09) (N = 2668)
- 6 (27.3%) of the 22 family members had committed suicide during the previous year
- Taking into account the average number of family members per household of 5.31 and the number of 2668 respondents that answered this question this produces a rate of 42.35 suicides per 100000 per year 2011

Alcohol problems

1) Alcohol problems of family members

- 9.2% of respondents reported that they had family members having "problems with drinking too much alcohol" (M = .10, SD = .32, N = 2678)
- 8.8% had one family member (236 persons), 0.3% two family members (16 persons), 0.1% three family members (9 persons)
- 48.2% of these drank heavily during the previous week, 69.4% during the previous months and 82.9% during the previous year

2) Alcohol consumption of respondents

The respondents' answers to their own alcohol consumption are shown in the following graphs. 59.1% of the respondents who reported consuming alcohol said they drink wine, 40.9% beer.




Fig. 13: Percentage of respondents consuming alcohol in given time periods (N = 2686)

- 86% of respondents said to drink less than monthly
- 9.5% of respondents said to drink regularly at least weekly




Fig. 14: Percentage of respondents consuming the given amount of alcohol per occasion (N = 922)

- Of the respondents consuming alcohol 71.2% drank less than half a liter or two cans per occasion
- 8.4% of them drank more than one liter or 5 cans per occasion
- 22.4% of respondents said that their alcohol consumption was causing problems with family, friends or their health
- An average of .25 respondents (25,3%) (SD = .435) reported that their drinking was causing financial problems (N = 922)
- According to the definition described in the methodology 1 % of all respondents are likely to suffer from alcohol addiction and 0.4% from alcohol abuse according to their own answers
- 4.0% of male respondents (M = .040, SD = .196, N = 873) versus 0.2% of female respondents (M = .002, SD = .041 N = 1813) are likely to suffer from alcohol problems (abuse or addiction)

Drugs

- 0.5% of the respondents (M = .01, SD = .119, N = 2675) reported they had one to five family members who were having a problem with taking drugs 28.6% of these during the previous month
- 3 (0.1%) of the respondents reported taking drugs daily and 2 (0.1%) less than three monthly, 99.8% reported never to take drugs (N = 2686)
- All the respondents consuming drugs were male (0.6% of male respondents (N = 872))
- 80% of these reported this caused rarely or sometimes problems with their families (N = 5)
- 80% of these reported that their drug consumption was causing financial problems (N = 5)

Gambling

- 1.7% of respondents (M = .02, SD = .175, N = 2675) had one to four family members with a gambling problem
- 30.4% of these had gambled during the previous week, 47.8% during the previous month and 82.6% within the previous year (N = 46)
- 4.8% of respondents gambled at least sometimes, 6.6% of the male respondents (N = 870) and 3.9% of the female respondents (N = 1809)
- 6.1% of respondents reported that their gambling habits caused many or some problems
- In an additional question 40.8% of respondents reported that their gambling was causing financial problems (N = 125)

For the frequency of gambling habits among the respondents and the results to the question: "Does your gambling ever cause any <u>problems</u>?" please see appendix in full report.

Cross-tables for `suicide attempts´, `in cage or chained´ and `alcohol problems´ with demographic variables

The following cross-table shows the number and percentage of respondents having an alcohol problem, having attempted suicide and having been chained or put in a cage by sex and urban or rural location.

		Suicide attempt In cage or chair		or chained	alcohol p	roblems	
		`	Yes	Yes		Yes	
		N	%	N	%	N	%
Male	Within male	15	1.7**	5	0.6	35	4.0**
	Within problem category		13.0**		62.5		92.1**
Female	Within female population	100	5.5**	3	0.2	3	0.2**
	Within problem category		87.0**		37.5		7.9**
Urban	Within urban population	44	5.1	3	0.3	9	1.0
	Within problem category		38.3		37.5		23.7
Rural	Within rural population	71	3.9	5	0.3	29	1.6
	Within problem category		61.7		62.5		76.3

Table7: Cross-table for respondents' alcohol problems, suicide attempts and being held captive in a cage or on chains and sex and living area

- Men had a nearly 20 times higher percentage of alcohol problems than women (highly significant)
- Women had a more than three times higher percentage of suicide attempts (highly significant)
- Males were three times more often affected with being chained or put in a cage due to a mental disorder (nearly significant)
- Acohol problems were more frequent in rural areas, whereas a higher percentage of the urban (5.1%) thatn the rural (3.9%) population tried to commit suicide (not significant)

Correlations between suicide attempts, in cage or chained and alcohol problems with demographic variables (age, years of education and economic situation)

- 76.3% of respondents with a probable alcohol problem had no savings or small debts. Alcohol
 problems correlated significantly negatively with savings per capita, meaning that the less wealthy
 suffered more from alcohol problems or vice versa alcohol problems lead to more spendings (Mann
 Whitney U: <.05)
- Alcohol problems correlated significantly negatively with age, meaning that the younger population was more affected. 2% of the group of 36 years or younger versus 1.1% of the group aged above 36 were affected
- Suicide attempts overall were more prominent in the less educated population (not significant)
- 87% of the respondents that had tried to commit suicide had no savings or were in debts (significant negative correlation between savings per capita and suicide attempts Mann Whitney U: <.05)
- Being chained or put in a cage of respondents correlated significantly negatively with savings per capita (Mann Whitney U: <.05) meaning that severe mental illness was associated with poverty

Summary: Major findings from the screening questionnaire

- Children problems were highest regarding aggressiveness (11.5% of respondents households with own children, 5.9% of respondents children)
- 8.8% of the respondents had experienced child abuse, described as physical or sexual violence or severe neglect (10.6% of male and 8.0% of female respondents)
- In 4.0% of the households there were one to two members affected with a probable schizophrenic or schizoaffective disorder
- About 0.74% of the family members were affected with a possible schizophreniform disorder (lifelong) and 0.2% of those were acutely affected with it
- Males were three times more affected with being chained or put in a cage due to a mental disorder
- 56.1% of respondents reported feeling irritable or angry, 60.5% out of these within the previous month
- 4.3% of the respondents had previously attempted to commit suicide (1.7% of male and 5.5% of female respondents) 22.6% of these within the previous month
- Respondents that had attempted to commit suicide suffer from much higher levels of psychological symptoms
- Women had a more than threefold rate of suicide attempts
- Suicide attempts overall were more prominent in the less educated population
- 87% of the respondents that had tried to commit suicide had no saving or were in debts, the poorer the respondents were the more they committed suicide attempts (significant)
- Alcohol problems were more frequent in rural areas, whereas suicide attempts occurred more
 often within the urban population
- In 0.8% of the households a member had committed suicide
- The suicide rate estimated for the general population was 42.35 suicides per 100000 per year (2010/2011)
- 9.2% of respondents had one to three family members with an alcohol problem
- 9.5% of respondents reported drinking regularly daily or weekly
- 25,3% of respondents consuming alcohol reported that their drinking was causing financial problems
- Men had a more than 4 fold rate of alcohol problems than women
- According to the definition described in the methodology 4.0% of male respondents versus 0.2% of female respondents are likely to suffer from alcohol problems (abuse or addiction)

III Traumatic experiences

Traumatic experiences during Civil War

- 64.1 % (1722) of the respondents had experienced the Civil War period comprising trauma during and shortly before and after the KR era out of these
- Only 7.7% had experienced or witnessed no traumatic events during CW
- The average number of traumatic events experienced was 7 (SD = 4.22, Max = 20)

The following graph shows the frequencies of different traumatic events experienced during the CW.




Fig. 15: Traumatic events experienced or witnessed by respondents during Civil War in % (N = 1714-1718)

The most and least frequently experienced traumatic events (in order of frequency)

Traumatic event	Experienced by	Witnessed by
Lack of food and water	75.1%	5.4%
Forced separation from family/friends	67.5%	6.1%
III health	46.3%	15.3%
Near death	46%	10.1%
Other traumatic events not in the list	18.2%	2.5%
Forced marriage	1.9%	1.6%
Sexual abuse/rape	0.6%	2.1%

Table 8: Frequencies of most and least experienced traumatic events

- Most frequently experienced and witnessed: lack of food and water (80.5), forced separation from family or friends (73.6 %), ill health (61.6 %), and being close to death (56.1%)
- Least frequently experienced and witnessed: forced marriage and sexual abuse or rape
- 18.2 % experienced and 2.5 % witnessed another traumatic event during CW not mentioned in the list

Traumatic experiences after Civil War

- Traumatic events after the CW were experienced or witnessed by 70.2% of the respondents
- 35.5% of respondents had experienced one, 19.5% two, and 8.7% three such events
- M = 1.31, SD = 1.28, Max = 9

The following graph shows the percentage of respondents having experienced or witnessed a traumatic event after the Civil War.




Fig16: Traumatic events experienced or witnessed by respondents after Civil War in % (N = 2687-2689)

- The three most frequent events were: fire, flood or other natural disasters experienced/witnessed by 53.4%, a life threatening accident by 25.2% and the witnessing of a bad injury or killing by 20.7%
- 13.9 % of respondents experienced/witnessed other extremely stressful or upsetting events
- Gang rape was not mentioned

Other stressful experiences



- 94.4 % of the respondents had experienced one to 12 stressful experiences
- 27.1 % of the respondents experienced two stressful events, 23.3% three and 15.9% four
- M = 2.93, SD =1.66, Max = 12

The following graph shows the percentages of the respondents' other stressful experiences.




Fig.17: Other difficult events experienced by respondents in % (N= 2668-2689)

- The five most frequently experienced events were financial worries, worries about problems of other family members, death of parents, grave illness of parents and worries about land mines
- Additionally 10.8 % of respondents worried about other problems not mentioned in the list

Traumatic events combined

- 87.9% of all respondents experienced or witnessed one to 24 traumatic events during and/ or after the CW (M = 5.8, SD = 5.11). 98.7% of all respondents had one to 31 traumatic and/or stressful experiences (M = 8.7, SD = 5.75)
- Traumatic events after CW were experienced more by men (M = 1.6 versus 1.2), as well as traumatic events during CW (M = 7.7 versus M = 6.6), whereas other stressful events were experienced slightly more frequently by women (M = 3.0 versus M = 2.9)

Correlations between the different traumatic/stressful experiences

Table 9 reflects relationships between different traumatic and stressful experiences at certain time periods. In the analysis, the amounts of traumatic events during and after the CW were also looked at separately as 'experienced' or 'witnessed' events. However, this did not show stronger correlations related to symptoms. Therefore, these correlations will not be shown in the results.

For the total amount of traumatic experiences during or after the CW the results for experienced and witnessed events were therefore combined as this also reflects the definition of a traumatic event according to DSM.

The following table shows inter-correlations between the sums of traumatic or other stressful life experiences at different times.

Different traumatic events experienced or witnessed							
Time Period	iod During Civil After Civil Other stressful						
	War	War	experiences				
During Civil War	1	-	-				
After Civil War	.274**	1	-				
Other stressful life	.217**	.227**	1				
experiences							

Table 9: Inter-correlations between various traumatic experiences ** = p< 0.001

The results of table 9 show that:

• There is a relationship between the number of traumatic experiences at different times. This suggests that those who have experienced more trauma during CW have also experienced more trauma after the war as well as other stressful life experiences

Summary: Traumatic experiences

• The most frequently experienced traumatic/stressful events:

During CW: lack of food and water, forced separation from family or friends, ill health and being close to death

After CW: fire, flood or other natural disaster, a life threatening accident and the witnessing of a bad injury or killing

Other: financial worries, worries about problems of other family members, death of parents, grave illness of parents and worries about land mines

- 70.2% of the respondents had experienced at least one traumatic event after CW
- 94.4 % of the respondents had experienced at least one of the stressful experiences
- The amount of other stressful life experiences such as poverty, death/illness in families, and marital problems has a high correlation with traumatic events experienced during and after the CW period

IV Self-reporting Symptom Scales

Harvard Trauma Questionnaire Symptoms

The following table shows the means (M), standard deviations (SD) and confidence intervals (CI) of the results from HTQ-Symptom scale for the total sample and for the male and female subsamples.


	HTQ Symptoms			% probable PTSD diagnosis		
	M	SD	95% CI		95% CI	
total	1.43	.40	1.41-1.44	2.7	2.1-3.3	
N = 2689						
women	1.46	.42	1.44-1.48	3.1	2.3-4.0	
N = 1813						
Men	1.36	.34	1.34-1.38	1.6	0.8-2.4	
N = 872						
= 36 years</td <td>1.37</td> <td>.38</td> <td>1.35-1.40</td> <td>1.5</td> <td>0.7-2.3</td>	1.37	.38	1.35-1.40	1.5	0.7-2.3	
N = 922						
> 36 years	1.45	.41	1.43-1.47	3.3	2.5-4.1	
N = 1767						

Table 10: M, SD and CI for HTQ scores and percentage of probable PTSD diagnosis

- The highest mean was found for the females
- 2.7% of the total respondents had HTQ-Symptoms scores above 2.5 the threshold score for probable diagnosis of PTSD (3.1 % of female and 1.6% of male respondents)

For the HTQ-Symptoms experienced by respondents above the total HTQ-Symptoms-mean of 1.43 please see graph in full report.

The following graph shows HTQ-Symptoms experienced by 9% or more of the total respondents as quite a bit or extremely.



- 1. Trouble sleeping
- 2. Avoiding activities that remind you of traumatic or hurtful events
- 3. Avoiding thoughts or feelings associated with traumatic or hurtful experiences
- 4. Feeling irritable
- 5. Difficulty performing work or daily tasks
- 6. Recurrent thought or memories of the most hurtful or terrifying events
- 7. Feeling jumpy, easily startled
- 8. Hopelessness
- 9. Feeling as if you don't have a future
- 10. Sudden emotional or physical reaction when reminded of the most hurtful or traumatic events

Fig.18: HTQ-Symptoms experienced quite a bit or extremely by >9% of respondents (N = 2681-2689)

- Trouble sleeping was the most common symptom experienced by 16.4% of respondents who replied it was a problem "quite a bit" or "extremely"
- The second and third most frequently experienced symptoms were avoidance symptoms with avoidance of activities reminding of traumatic experiences with 13.7% of respondents selecting quite a bit or extremely and avoidance of thoughts or feelings associated with traumatic experiences (13.3% of respondents)
- Feelings of emotional numbing (3.1%), shame (5.1%) and dissociative experiences (loss of memory (4.9%), feeling detached (4.6%), feeling split (5%)) were least experienced

Cambodian Addendum of Culturally Sensitive Items (now: C-SSI)



The authors divided this scale into two subscales according to Hinton (2011 c): the subscale of syndromes comprises the six items `thinking too much' kit chraeun, `weak heart' khsoy besdong, `wind hitting up from your stomach' khyal chral leung leur pi krapas thveur ouy nak khlarch stas, `wind attacks' gaeut khyal, `numb arms and legs' cok day cog ceung and `sleep paralysis' khmaoch songot, which was weighed double, because of its rareness and extreme stress associated as suggested by Hinton (2011 c, personal note). This leads to a total maximum score of 4.29 instead of 4. The subscale of symptoms comprises the following eight items: `weakness' khsoy, `standing and feeling dizzy' krokchhor heuy wul muk theang thong, `dizziness' wul muk, `tinnitus' hung treujeu, `neck soreness' rooy go, `blurry vision' prul phaneik and `poor appetite' mun khlieun.

The following table shows the M, SD and 95%CI of the C-SSI total and its subscales.

	M	SD	95% CI
C-SSI total	1.76	0.50	1.75-1.78
N = 2689			
C-SSI syndromes	1.86	0.51	1.84-1.88
C-SSI symptoms	1.69	0.54	1.67-1.71
C-SSI total women	1.81	.51	1.78-1.83
N = 1813			
C-SSI total men	1.67	.44	1.64-1.70
N = 872			

Table 11: M, SD and Cl of C-SSI total, C-SSI syndromes, C-SSI symptoms

For the items of the C-SSI that were most prominent among the respondents of this study (M > total M of 1.76) please see full version of report.

The following graph shows the C-SSI symptoms which were reported as occurring "extremely" or "quite a bit" by over 15 % of the respondents.




Fig.19: C-SSI items experienced extremely or quite a bit by > 15% of the respondents (N = 2683-2689)

- The three items experienced most frequently by the respondents were the syndromes of `thinking
 too much' with 37.4% and of `numb arms and legs' with 24.5% and the symptom of `blurry vision'
 with 22.8%
- `Sleep paralysis' (2.9%), `poor appetite' (6.4%) and `cold hands and feet' (M = 10.4%) were the least frequently experienced items

The Hopkins Symptom Checklist – 25 for Anxiety and Depression (HSCL-25)

The following tables show the means (M), standard deviations (SD) and confidence intervals (CI) of the results from HSCL-25 for anxiety and depression scores and percentages of probable diagnosis for the total sample and for the male and female sub-samples. The highest possible score is 4.

	Anxiety			% probable diagnosis of anxiety		
	M	SD	95% CI	M	95% CI	
total	1.51	.47	1.50-1.53	27.4	25.7-29.1	
N = 2689						
Women	1.57	.49	1.55-1.59	31.7	29.6-33.9	
N = 1813						
Men	1.40	.39	1.37-1.42	18.4	15.8-21.0	
N = 872						
= 36 years</td <td>1.43</td> <td>.43</td> <td>1.41-1.46</td> <td>20.9</td> <td>18.3-23.5</td>	1.43	.43	1.41-1.46	20.9	18.3-23.5	
N = 923						
> 36 years	1.56	.48	1.53-1.58	30.8	28.7-33.0	
N = 1766						

Table 12: M, SD and CI for HSCL-25 scores of anxiety

	Depression			% probable diagnosis of depression		
	M	SD	95% CI	M	95% CI	
total	1.44	.39	1.43-1.46	16.7	15.3-18.1	
N = 2689						
Women	1.48	.41	1.46-1.50	19.7	17.9-21.5	
N = 1813						
Men	1.36	.31	1.34-1.38	10.2	8.2-12.2	
N = 872						
= 36 years</td <td>1.38</td> <td>.37</td> <td>1.36-1.41</td> <td>12.9</td> <td>10.7-15.1</td>	1.38	.37	1.36-1.41	12.9	10.7-15.1	
N = 923						
> 36 years	1.47	.40	1.46-1.49	18.6	16.8-20.5	
N = 1766						

Table 13: M, SD and CI for HSCL-25 scores of depression

- Considering the cut-off score of a mean of 1.75 for the probable diagnosis of anxiety and depression respectively, 27.4% of the respondents suffered from anxiety (95% CI: 25.7%- 29.1%) and 16.7% from depression (95% CI: 15.3 %- 18.1%)
- The percentage for women was 31.7% for probable anxiety, 19.7% for probable depression and for men 18.4% and 10.2% respectively

1) Symptoms of anxiety

For the most prominent symptoms of anxiety of or above the total mean (M > 1.51) please see full version of report.

The following graph shows the HSCL-25 symptoms of anxiety as experienced extremely or quite a bit by more than 10% of the respondents.




Fig. 20: HSCL-25 symptoms of anxiety experienced extremely or quite a bit by > 10% of respondents (N = 2687-2689)

 The most common symptom was headache, followed by faintness/dizziness or weakness, and heart-pounding

2) Symptoms of depression

For the most prominent symptoms of depression above the total mean (M > 1.44) please see full version of report.

The following graph shows the HSCL-25 symptoms of depression as experienced extremely or quite a bit by more than 10% of the respondents.




Fig. 21: HSCL-25 symptoms of depression experienced extremely or quite a bit by > 10% of respondents (N = 2686-2689)

- The three most frequently reported symptoms are worrying too much, difficulty falling or staying asleep, and low energy/apathy
- Additionally, loss of sexual interest is the symptom which is experienced 'extremely' by the highest percentage of respondents (5.7%) out of all depressive symptoms
- The thought of ending one's life (0.9%), feeling no interest in things (3.2%) and the feeling of being trapped or caught (3.7%) were experienced least frequently

Summary of findings of symptoms experienced by respondents

	М	SD	Max	Maximum score possible	% above threshold score for diagnosis
HTQ-Symptoms	1.43	.40	3.75	4	2.7
HSCL-25 Anxiety	1.51	.47	3.8	4	27.4
HSCL-25 Depression	1.44	.39	3.8	4	16.7
C-SSI Total	1.76	.50	4.29	4.29	

Symptoms experienced most/least frequently:

- HTQ: most: trouble sleeping, feeling irritable, avoidance of activities and thoughts associated with traumatic experiences, difficulty performing work or daily tasks; least: feelings of emotional numbing, shame, loss of memory, feeling split
- C-SSI: most: thinking too much, numb arms and legs, standing and feeling dizzy, blurry vision; least: sleep paralysis, poor appetite, cold hands and feet
- HSCL anxiety: headache, faintness/dizziness or weakness, heart-pounding
- HSCL depression: worrying too much, difficulty falling or staying asleep, low energy/apathy

V Mental health and demographic factors

Gender

The following graph shows the different symptom scale mean scores by sex of respondents.




Fig. 22: Mean total scores of CSSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by sex

 Clearly higher mean scores for all scales can be seen for female in comparison to male respondents on a highly significant level (Mann Whitney U: p<.001)

Age

The following graph shows the different symptom scale means by age below or above 36.




Fig.23: Mean total scores of CSSI HSCL-25 Anxiety and Depression and HTQ-Symptoms by age split

 Higher age is related to higher scores in all the symptom scales with high significance for all scales (Mann Whitney U: p<.001)

Marital status

The following graph shows the different symptom scale means by marital status.




Fig. 24: Mean total scores of C-SSI HSCL-25 Anxiety and Depression and HTQ-Symptoms by marital status

- Single persons have the lowest scores in all symptom scales and widowed persons mainly the highest
- Divorced respondents have lower scores for somatic symptoms (C-SSI) than married ones, whereas this is opposite for probable anxiety, depression and PTSD
- The differences are all highly significant (Kruskal Wallis: p<.001)

Location

The following graph shows the relation between the different symptom scale means and the location urban or rural.




Fig. 25: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by location

• The urban population suffers a bit more from traumatic symptoms and depression whereas the rural population suffers more from anxiety and typical Cambodian and bodily symptoms. This is only significant for C-SSI and HTQ (Mann Whitney U: p<0.05).

Education

The following graph shows the different symptom scale means by years of education per respondent in categories of no, primary, secondary, high school and university education.

Scale means by education

mean no 2 1.86 primary 1.8 1.67 1.54 secondary 1.5 1.6 1.44 high school 1.4 university 1.2 1.57 0.8 1.36 0.6 0.4 0.2 c-ssi HSCL-25 HSCL-25 HTQ Depression Symptoms

Fig. 26: Mean total scores of C-SSI, HSCL-25 Anxiety and Depression and HTQ-Symptoms by education in categories

- Overall respondents with no or primary education reach/have highest scores in all of the symptom scales
- The highest differences were found for C-SSI and anxiety, the least for HTQ
- The differences were all highly significant (Kruskal Wallis: p<.001)

Economic situation

The following graph shows the different symptom scale means by average savings per respondent in category midpoints.




Fig. 27: Mean total scores of C-SSI, HSCL-25 and HTQ-Symptoms by savings per person in the respondents' household

- A clear decline of the C-SSI, HSCL-25 anxiety and depression and HTQ-Symptom scores can be seen with rising savings
- This is highly significant for Anxiety and Depression (Kruskal Wallis: p<0.01) and significant for C-SSI (p<.05)

Correlations between different symptom scores and demographic factors

The following table shows correlations between mean syndrome scores and demographic factors such

Mean total scores of				
	HSCL-25 Anxiety	HSCL-25 Depression	C-SSI Total	HTQ Symptoms
Age	.152**	.155**	.267**	.123**
Years of education	161**	130**	209**	063**
Savings per capita	083**	106**	059**	078**

Table 14: Spearman correlations between mean symptom scale scores and demographic factors with significance level ** <0.001

The results of table 14 show that:

- All of these demographic variables correlated highly significantly with all of the syndrome scales
- Compared with HSCL-25 and C-SSI the HTQ-Symptoms showed the smallest significant correlations with demographic data variables. The smallest association was found with level of education (-.063). This indicated that higher educated respondents were slightly less frequently traumatized
- The highest correlations with age and years of education were reached by C-SSI. This means that these syndromes and symptoms increase most with higher age and lower education (negative correlation). C-SSI is least associated with the economic status
- The HSCL-25 Anxiety scale had the second highest correlation with years of education. This
 means that education has an important influence on the development of anxiety or vice versa
 anxiety hinders educational progress
- The HSCL-25 Depression scale correlated highest of all scales with savings per capita meaning that the more depressed a person is the less it can earn or vice versa the less it earns the more

Summary: Symptom scales and demographic factors

- Compared with HSCL-25 and C-SSI the HTQ- Symptoms showed the smallest significant correlations with demographic data variables such as age, sex and years of education
- C-SSI scores increase most with higher age and lower education, anxiety has the second highest association with these
- Gender-dependence is high for all of the scales with significantly higher proportions of women suffering from psychological symptoms
- Increasing poverty is associated highest with increasing depressive symptoms
- Single respondents suffer least, widowed respondents most from psychological symptoms
- The urban population suffers significantly more from trauma-symptoms, the rural population more from typical cultural syndromes

VI Services

Service utilization, costs and knowledge

The following graph shows percentage of yes-answers to the following questions: "Is anybody providing help for the above problems?" "Is it affordable?" and "Do you know about psychology?"




Fig. 28: Percentage of yes answers to questions about services economics and knowledge

- Only 24.1% of the respondents stated that someone was providing help for them if one of the problems mentioned during the interview occurred
- Of the ones that/who received treatment 61.4% reported it was affordable, but only 132 respondents answered that question
- 19% of respondents claimed to know something about psychology, 29.9% of male and 13.6% of female respondents

The following graph shows the percentage of respondents that/who approached the mentioned sources for help. Multiple responses were possible.




Fig. 29: Percentage of respondents who approached the mentioned sources for help (N = 2688-2689)

- At the top of services approached are low level medical services such as health posts, pharmacies and small clinics (probably respondents understood also small drug stores by this), the second important resource approached for help by 50.7% were the respondents' families
- 20.7 27.9% of all respondent contacted traditional spiritual services such as monks, mediums or traditional healers for help
- Doctors and neighbors were contacted least (3.9 and 0.4% respectively)
- 13.5% reported they approached other sources for help
- There were no major differences for gender in seeking help behavior

Needs for services

- After an explanation about what psychology means, 7.6% of respondents answered with yes to the
 question "Are there any psychological programs or NGOs providing help in this field near your
 village?" (N = 2028).
- To the question "Would you like to have a psychological service at your health center" 97.2% answered with "yes", 0.5% with "maybe" and 2.3% with "no" (N = 2658).
- 35.7% of respondents stated that they would contact a psychologist for their problems related to the interview (N = 2689).



Overall the economic situation was poor for the survey respondents. Over 48% (48.2%) of the households earned less than 50 USD per month which is nearly equal to the findings by Sonis et al. (2009). Furthermore 87.4% had either no savings or were in debt. In addition, numerous issues related to mental health were apparent in the sample.

I Screening questionnaire Children problems

High rates of problems in children were found such as high rates of aggression. These high rates may be interpreted as an indicator for second generation affects from the traumatic past experiences as they are linked with high reports of aggressiveness of respondents and other family members. Several difficulties were met when trying to interpret these findings. There are few reports on prevalence rates for disorders in childhood and adolescence. Thus prevalence figures vary considerably, e.g. 12.8% in India for children aged 1 to 16 and 22.5% in Switzerland for children aged 1 to 15 (WHO, 2001). The questions in this study focused on indirect indicators for children's problems and didn't cover the whole spectrum of disorders. Additionally, it is difficult to draw clear borders between normal development and developmental pathologies. The terms "aggressive behavior" and problems" weren't specified, so it remains open what respondents understood by these terms. Furthermore, taking into consideration that many interviews lacked sufficient privacy it may be that some cases remained concealed and the real numbers are even higher. However even with these uncertainties it can be concluded from these results that family and school counseling as well as psychosocial services for children and adolescents should become a high priority in mental health planning in Cambodia as this generation is in great need and will determine Cambodia's future.

Traumatic events

Over 8% of the respondents reported that they had been abused as a child (8.8%), and 4% of respondents reported that at least one of their family members had been victims of child abuse. Because many of the interviews lacked privacy (approximately 50%) the real numbers may again be higher. The discrepancy between the incidence rates reported by the respondents referring to themselves as opposed to other family members also suggests a taboo of this topic in the families. The high figures are in line with results from the Stop Violence Against Us- Report (Miles, 2006 a) that revealed that nearly a quarter of those interviewed witnessed the rape of a child in their community. A study from the US revealed incidence rates of 10.2% of maltreated children and 6.1% of sexual assault victims among 0 to 17 year olds (Finkelhor, Turner, Ormrod & Hamby, 2009). Contrary to the very common view of childhood abuse as a major problem for girls, the figures from the present study are higher for men (10.6%) than women (8.0%). Some organizations in Cambodia do try to address this fact of abuse of boys. Survey results from the Stop Violence Against Us- Report part II (Miles, 2006b) revealed that 13.3 percent of girls admitted having been sexually touched on the genitals before reaching the age of nine years as opposed to 15.7 % of boys. The figures found in the present survey are alarming as the impact of child abuse is undoubtedly one of the most detrimental to the human psyche and can lead to psychological problems such many depression, anxiety, addictive disorders and suicidal tendency in the long term (van der Kolk, Hoppner & Crozier, 2001 Felitti et al., 1998) as well as pose a great socioeconomic burden (Fang, Brown, Florence & Mercy, 2012).

The reported numbers for trafficking in this report were 0.4% of respondents and 0.7% of family

members, which is again quite high in spite of the lack of privacy during the interviews. According to a report from the MoWA (2008), incidents of rape and sexual assault are increasing (MoWA, 2008). In an international survey conducted by the National Criminal Justice Reference Service (2011) the prevalence of trafficking in Asia was estimated at 3/1000 inhabitants as compared to 1.8/1000 inhabitants worldwide.

However numbers for rape lay with 0.4% of respondents and their family members which is far below the U.S. rate of 14.8% life time experience of rape for women (Tjaden & Thoennes, 2000). This may be in part due to a higher rate of violence in the U.S., but it probably results from the fact that this issue is an especially sensitive one and probably many respondents feared stigmatization and negative consequences and therefore withheld information.

Probable psychotic/ schizoaffective/ bipolar disorders

This study provides the first data about probable schizoaffective psychotic or disorders Cambodia although the data are indirect in nature. We calculated in the present data a lifetime prevalence for probable psychotic disorders of 0.76% for the total family members and a point prevalence of about 0.2%. The higher proportion of respondents that experienced being put in cages or chained because of a mental problem themselves in comparison with their family members seems difficult to explain as such events are quite dramatic and difficult to forget. Possibly a fear of offending family members or of getting in trouble for having violated human rights prevented more information about these cases. Compared to data from WHO (2001) reporting a point prevalence for schizophrenia of 0.4% (GBD, 2000) and a lifelong prevalence of under 2% for schizophrenia and bipolar disorders the results from this study seem guite realistic. Schizophrenic and bipolar disorders are chronic disorders, which means that they have high relapse rates and need continuous medication with surveillance via regular follow-up consultations as compliance and insight into being ill is generally very low. Therefore these types of disorders account for 18.5% of the cases presenting at governmental mental health services although their prevalence is much lower

(Stewart et al., 2010). Considering the high need for regular treatment and the very low availability mental health services especially Cambodia's rural areas (McLaughlin & Wickeri 2012) the above described facts display an alarming outcome of the present study. Patients with these kinds of disorders are often forced to travel to Phnom Penh for more adequate treatment, which is a considerable financial burden for the family. In addition to the high expenses the families have perhaps spent already on inadequate traditional treatments and other treatment costs they have to deal with travel expenses and often with missing income from perhaps two potential income generators (the patient and caregiver). Many families have to sell their land and property or borrow money which they often can't pay back (a potential reason for suicides). The relationship between poverty and probable psychotic disorders was confirmed in this study and may be due to a shift from the wealthier to the poorer population following the onset of this disorder as described in other countries (drift hypothesis; Buck & Morrison, 1988). The often unavoidable consequence of putting family members affected with these disorders in cages or chaining them due to insufficient financial resources is a great risk for further psychological harm and traumatization, which often may augment and prolong the problem, as well as it being a serious violation of the patient's human rights (see also McLaughlin & Wickeri, 2012; Stewart et al., 2010). Cambodia is a party in the International Covenant on Economic Social and Cultural Rights on "the right [...] to the [....] highest attainable standard of physical and mental health." (ICESCR, 1996) and has guaranteed free medical consultations for the poor and full consideration of disease prevention and medical treatment according to article 73 of the constitution (1993). However, Cambodia is far from upholding these promises and a higher allocation of money to the mental health sector is urgently required in order to meet the needs of acutely and chronically mentally ill. In addition mental health literacy must be improved in order to enable the population to react more adequately these disorders and to promptly find appropriate help.

Suicidal tendency

Another very alarming result are the high rates of attempted and committed suicides. The rate of suicide attempts for the respondents themselves was at 4.2% (of which 44.3% was within the previous year), for family members at 0.8% (50% within previous year). The discrepancy between the rate for the respondents themselves and their family members may reflect the lack of communication and the large amount of unnoticed suicide attempts in the families. To the researchers knowledge many clients attempt suicide without their families realizing. In 0.8% of the 2690 households interviewed in this study at least one family member had successfully committed suicide. Of these suicides, 27.3% had occurred during the past year which produces a rate of 42.35 suicides /100,000 population per year (including both males and females). The Cambodian suicide rates therefore exceed worldwide numbers more than twofold with an average of 16 suicides per 100,000 people per year (WHO, 2011a). A study from Thailand (Lotrakul, 2006) found that between 1998 and 2003 there was an average suicide rate of 7.9/100,000 and a male to female ratio of 3.4:1 with males aged 25-29 at the highest rate. Compared with this culturally close neighbor, the Cambodian rate seems especially high as it exceeds the Thai numbers more than fivefold. Taking into account the unclear distribution, the figures found in this study come close to the highest numbers seen around the globe. For comparison, a rate of 74.6/100000 has been reported for the male population in Lithuania and 63.6/100,000 for the male population in Belarus from 2003 (WHO, 2011c).

In comparison with the rates for accomplished suicide, the numbers for attempted suicides seem even higher as the ratio is higher (up to 50fold) than the data reported by WHO (2011a) which is 20fold.

Regarding the gender distribution for attempted suicides the present study found 1.7% of men and 5.5% of women. In a study by Jegannathan & Gunnar (2011) who interviewed 320 students from two high-schools in Takhmau suicidal plans were reported more often by teenage boys than teenage girls (M = 17.3%, F = 5.6%), whereas girls reported more attempts (M = 0.6%, F = 7.8%).

The outcome confirms the researchers' expectation and clinical experience. international trend of increased suicidal behaviors in younger people and the higher percentage of women attempting to commit suicide (WHO, 2011 a) are confirmed in this study (>3:1 female: male) on a significant level. Cambodian youth seem to be at the highest risk for suicidal behavior. Frequently the younger generation has the duty and burden to earn the family's income and to support other family members. Also a strong fear of burdening the family seems to drive many to consider suicide. One of the most commonly mentioned worries in this study was the lack of money. In fact, the family savings per capita was negatively correlated with the rate of suicide attempts.

Suicidal tendency also correlated highly with the rate of psychopathology as measured by the HSCL-25 Anxiety and Depression scale, the HTQ -Symptoms and CSSI scales. The loss of fitness for employment and of earning capacity due to mental illness strains the family, especially in the context of the ever-present poverty in Cambodia. Debts, pressure and worries increase and corner many in a seemingly intractable situation. These results indicate that poverty reduction, a better infrastructure for mental health and increasing services in this field could potentially reduce suicidal tendencies. One other factor with impact on suicidal tendency may be the number of HIV infections as suggested in the Thai study (Lotrakul, 2006).

The topic of suicide is also prevalent in music videos and movies broadcast on TV and shown all over Cambodia such as during bus trips. These often suggest suicide as an adequate reaction to lost love and abandonment in partnerships. More detailed research should be conducted on this topic, but it seems advisable to raise awareness on this issue and call for other entertainment. Suicide attempts also have to be considered as cries for help and attention which due to a low level of communication cannot be achieved in other ways. This can be understood as an appeal for the improvement of interpersonal communication skills and emotional sharing.

Addictive disorders

The figures of 1% point prevalence for probable alcohol addiction and 0.4% point prevalence for probable alcohol abuse by respondents in this study seem quite realistic in comparison with the GBD 2000 point prevalence for harmful use of alcohol or alcohol dependency at 1.7% overall (WHO, 2001), especially when taking into account the oversampling of female respondents in this study. The gender distribution was about four to five times as uneven as suggested by WHO figures (2-3:1 WHO, 2011c) with a ratio of 35 men to 3 women probably suffering from alcohol problems in the present study.

As these figures rely on self-report and alcohol-dependents tend to underreport, the actual number may well exceed this prevalence rate. At the same time it has to be taken into account that these figures are based on indirect indicators and not on clinical diagnostic interviews and examinations.

A worrying fact is that alcohol problems were reported nearly twice as frequently in the younger age group (<36) (2%) compared to the older age group (1.1%) and correlations showed significant association between alcohol problems and younger age as well as poverty. These results mirror findings from other countries. Correlations with poverty can be explained either by the "drift hypothesis" of increasing poverty following addictive disorders (Buck & Morrison, 1988) or by the "social causation thesis" of use of alcohol as self-medication to cope with increasing financial worries, trauma and depression (Perry, 1996). The poorest respondents however were less likely to suffer from alcohol problems probably due to a lack of financial resources.

In order to prevent a further increase of this problem, poverty should be reduced in Cambodia, and the Cambodia media should support psychoeducative programs on radio and TV.

Gambling is a problem as common as alcohol problems. Here the fact that the term "gambling" was not clearly defined complicates the interpretation of results, because many families interpret gambling more in terms of a social activity than an addictive behavior. Taking into account that 4.7% of respondents admitted they gamble and 40.8% of these stated that this .

gambling was causing financial problems it can be assumed that around 1.5% are likely to suffer from some degree of gambling addiction. Because of the self-reporting structure of the interview fear of stigmatization may have reduced this prevalence rate.

A study by Marshall, Elliott and Schell (2009) of 127 Cambodian refugees in the USA reported that 13.9% of the participants met their criteria for "disordered gambling". Being male, being married, and being exposed to high levels of trauma were variables associated significantly with gambling. The prevalence of gambling may have increased in this refugee population because of "uprooting" and the stress of adapting to a new country and culture.

The problem of gambling is especially important to notice as the danger of impoverishment caused by gambling is a great risk for the whole family.

Drug consumption in comparison seems much less frequent in the Cambodian population. As no typical signs of addiction such as craving, withdrawal or tolerance were collected it is difficult to estimate prevalence rates. The most important findings here were that 0.1% of the respondents stated that they take drugs on a daily basis. Another 0.1% responded that they take drugs more frequently than every three months and 80% of these individuals had financial problems because of this.

Drug addiction often is more severe and difficult to treat than other addictive disorders and the problem seems to be much more prominent in urban areas, especially Phnom Penh. This problem has therefore obtained already great attention reflected in the new Mental Health and Substance Misuse Strategic Plan 2011-2015 (Ministry of Health, 2011). A substitution program has been set up at the Russian Hospital in collaboration with WHO and FHI. Counseling is added in cooperation with the DP.

Major outcomes and discussion of screening part

- The prevalence figures of difficult behaviors in children and adolescents seem high and sug gest that school counseling should be regularly introduced in Cambodia, however these figures are difficult to compare.
- Child abuse seems to be an outstanding/salient problem with men being more frequently affected, this is an alarming result considering its extremely detrimental impact.
- The problem of trafficking was found a little higher than suggested by previous data.
- Schizophrenic disorders and the like seem to occur at a fairly common rate, but pose a major
 problem regarding appropriate treatment facilities and human rights issues. 0.3% of respondents had experienced being chained or put in a cage themselves. Here improvement of mental health literacy in the population and intermediaries such as traditional healers and monks
 as well as increase of psychiatric out-patient clinics are major tasks.
- Suicidal tendency was found a major issue in the Cambodian society. The rate of 42.35/100000 encompasses worldwide figures more than 2.5fold. 4.3% of respondents had previously tried to commit suicide (44.3% during previous year), which is up to 50times the rate of accomplished suicides compared to a worldwide ration of 20:1. The international trend to younger people and the higher percentage of women attempting to commit suicide found in the literature are confirmed in this study (>3:1 female to male) on a significant level. A significant negative correation between the families' savings per capita and suicide attempts was found. This result clearly indicates that poverty reduction would probably prevent suicide at tempts most effectively. The highly significant association between suicide attempts and psy chopathology indicates that an improvement of general mental health care would probably al so lead to a significant reduction of suicidal tendencies.

II Traumatic experiences (Harvard Questionnaire)

As expected, traumatic events during the Civil War were experienced by a very large percentage of respondents above the age of 36 (92.3%).

In the following table results from this study are

compared with results from literature. Dubois et al. (2004) refer to lifelong traumatic events experienced in Kampong Cham. Sonis et al. (2009) to traumatic events experienced by a population over 35, and Mollica et al. (1993) to a refugee population.

Traumatic events experienced during	Present study	Study by Dubois et al. (2004)	Study by Sonis et al. (2009)	Study by Mollica et al. (1993)
Lack of food and water	75.1%	69.7%		
Forced separation from family	67.5%	18.9%	81.5%	
III health	46.3%	40.1%		
Close to death	46%	32.7%	50.1%	
Lost/kidnapped	34.9%	9.2%		
Totured	18.9%	8.3%	31%	36%
Murder of family/friend	8%	16.4%	22.1%	54%
Combat situation	9.7%	18.7%	62.8%	
Imprisonment	5.2%	4.4%	8.3%	

Table 15: Traumatic experiences during CW in comparison with other studies conducted in Cambodia

In total this study's findings seem guite consistent with other studies although the different study designs make the comparisons difficult. Still the findings in the present study for many events (torture and imprisonment for example) lie close to or in between previous findings. Greater differences are found for being lost/ kidnapped (higher percentage in the present study) and murder of family/ friends and combat situations (lower percentage in the present study). Further analysis by provinces can give further insight into the traumatic events experienced by Cambodians.

Also 70.2% of the total sample experienced

traumatic events after the Civil War, which seems very high. Natural disasters and life threatening accidents were experienced most. This calls for a more active preparation for trauma response not only on a pragmatic level of basic physical needs but also on a psychological level. As described in a master thesis by Sieng (2011) a well-trained team of psychologists and psychiatrists should be established for acute interventions in order to prevent a later onset of post- traumatic stress disorders as best as possible.

The following table compares the results obtained in the present study with the studies by Sonis et al. (2009) and Dubois (2004).

Traumatic experiences after the Civil War	Present study	Study by Sonis et al. Cohort < 35 years (2009)	Study by Dubois et al. (2004)	
Fire, flood, natural disaster	33.8%	48.9%		
Life threatening accident	16.0%	22.7%	16.9%	
Witnessed someone badly	15.7%	38.0%		
Seriously physically attacked	2.6%	5.8%		

Table 16:Traumatic experiences after CW in comparison with other studies conducted in Cambodia

The percentages seen in this study are 30 to 60% lower than results from the study by Sonis et al. (2009), but very close to that reported by Dubois et al. (2004).

The present study's findings regarding other stressful life experiences cannot be compared as none of the other studies collected these data. Alarming are especially the high amount of financial worries (86% of respondents), worries regarding insufficient health care (illnesses and deaths) as well as continuing high fears of landmines (15.9%). Dubois et al. (2004) noticed the high impact of poverty related issues on mental health as well. They distinguished between traumatic experiences related to poverty on one hand and mass violence on the other hand. Of their respondents, 55.7% experienced both types of events, 19.9% poverty alone, 11.1% violence alone and only 13.3% never experienced any such events.

The strong significant correlations among the various traumatic experiences at different times or in different intensity reflects the high amount of multiply traumatized individuals as well as the impact of traumatization on future vulnerability for

such traumatic experiences. Relatedly, previous research has shown that women that have been raped become victims of such crimes again more easily than others. This happens probably due to dissociative "freeze" reactions and lack of resilience (see also: Anda et al. 2006).

In a meta-analysis of 32 studies on violence and mental health in low and middle income countries by Wagner, Sergi, Cleusa, Martin & Jair de Jesus (2009), it was found that a substantial part of mental health problems can be attributed to violence. In the general population, the highest prevalence rates of PTSD were associated with sexual and domestic violence, kidnapping, and cumulative trauma exposure.

The correlations between the different experiences of traumatic and stressful events and the scores of the different symptom scales and regression analysis will be investigated further by the DP at RUPP.

III Symptom scales

Posttraumatic stress symptoms HTQ

The prevalence rate of probable PTSD for the total population aged 21 years and older of 2.7% in the present survey is 7 times higher than in the general population worldwide with 0.4% (WHO, 2001). The high rate of PTSD in the Cambodian sample is not surprising in light of the country's particularly horrific history. However it is much lower than in previous studies conducted with Cambodians:

 Carlson & Rosser 1991 refugees:
 86% (DES)

 Mollica et al. 1993 refugees:
 15% (HTQ)

 de Jong et al. 2003:
 28.4% (CIDI)

 Dubois et al. 2004:
 7.3% (HTQ)

Sonis et al. 2009: 11.2%(PTSD Checklist)

(oversampled for old age)

It is difficult to compare results with these studies because different diagnostic tools were used. In a comparative study across four countries, de Jong et al. (2001) found the following prevalence rates of PTSD: Algeria 37%, Cambodia 28%, Gaza 18%, Ethiopia 16%. These data have to be interpreted carefully, since the PTSD prevalence rates were estimated within very different time spans between the time when the actual traumatic events had occurred and the assessment of symptoms. This makes the data difficult to compare as PTSD usually develops and decreases in relation to time. In addition, in Algeria terrorist attacks were still ongoing during the data collection, so the diagnosis of PTSD may have been mixed up with acute stress reactions.

The relationship between age and PTSD (older age was associated with higher PTSD rates) found by de Jong, Komproe & Ommeren (2001) and Sonis et al. (2009) is consistent with findings from the West (Kessler, Sonnega, Bromet & Nelson, 1995) and was confirmed in the present study.

The prevalence rate Sonis et al. found 2009 is more than six times the one in the US (Field, 2011). In their study Sonis et al. chose a threshold score for PTSD of 44 instead of 50 as originally defined because it demonstrated a better balance of sensitivity and specificity according to other studies (Sonis et al., 2009, Brewin 2005).

Over all it can be seen that the trend is that the refugee populations suffer to a much higher

degree from PTSD and that the rates of PTSD decrease in Cambodia over time especially looking at the three studies using the HTQ-Symptoms scale. This is probably due to the fact that more individuals without the experience of the KR regime take part in the studies and maybe also that the prevalence of PTSD in traumatized individuals may decrease over time (McFarlane & Papay, 1994). Also the timing of the surveys may be an important element in the events reported by the traumatized subjects (Willis, 1998). Staff from Transcultural Psychosocial Organization (TPO) estimates the probable rate of PTSD at 10 to 18% based on their clinical experience at TPO's out-patient clinic (McLaughlin & Wickeri, 2012). This may be the result of a more clinical impression as presented by data from Sonis et al. (see above) and/or due to TPO's good reputation especially in the field of trauma which attracts cli-The rates found by Cambodia's ents. governmental psychiatric services are lower (reactions to severe stress and adjustment disorders 1.9% of the population visiting health care centers) (Stewart et al., 2010).

HTQ-Symptom scores were significantly higher in the female population. This corresponds to findings by de Jong et al. (2003) in Algeria and Cambodia. HTQ-Symptom scores significantly positively correlated with age (more PTSD with older age) and significantly negatively correlated with years of education (higher amount of education less vulnerable for PTSD), but with a weaker association than the other scales had. The relationships between age/ education and PTSD are consistent with findings from de Jong et al. (2001) and from the West (Kessler, Sonnega, Bromet & Nelson, 1995). HTQ-Symptoms were significantly more prominent in the urban population, which can possibly be explained by more traffic accidents and the greater burden of trauma from the KR on urban population.

With regard to the discussion about cultural differences in the expression of traumatic symptoms it was found in the present study that the second and third most experienced problems listed in the HTQ-Symptoms scale were avoidance of thoughts or feelings and activities. Re-experiencing and hyper-arousal symptoms only appeared on place six and seven (recurrent thoughts or memories, feeling easily startled).

The most prominent symptom of "trouble sleeping" can be interpreted as a sign of hyperarousal, re-experiencing or avoidance. High rates of sleeping difficulties were also seen in the HSCL -25 Depression scale. However findings from other studies differ to different degrees mainly depending on the scales used.

Present study: Avoidance > Reexperiencing > Hyperarousal

De Jong: Reexperiencing > Avoidance > Hyperarousal

Dubois: Reexperiencing > Avoidance

Sonis: Reexperiencing > Hyperarousal > Avoidance

A more detailed analysis of the different instruments, their clusters and outcomes is necessary to draw correct conclusions on this issue.

According to findings from Sonis et al. (2009) PTSD was associated with mental disability in a model controlled for demographic covariates.

Cambodian Symptom and Syndrome Inventory (C-SSI)

The mean average score in the present study for the C-SSI was 1.76 out of 4.29 with a significant gender difference. The most prominent symptom "thinking too much" was corresponding to the most prominent symptom of the HSCL-25 depression scale: worries. Other common ones were numb arms and legs, standing and feeling dizzy and blurry vision. Three of these belong to the subscale of cultural syndromes linked to a wide spectrum of cultural beliefs and interpretations, which can be regarded as a confirmation of the importance of these beliefs and associations (Hinton et al., 2005 b).

The C-SSI and HTQ-Symptoms scales were the only scales to show significant differences between the rural and urban population. The rural

population suffered more from cultural somatic symptoms probably due to their more traditional roots and beliefs.

Anxiety and depression (HSCL-25)

The prevalence rates for acute anxiety and depressive disorders measured with the HSCL-25 for the 2690 respondents were high:

Anxiety: total: 27.4% men: 18.4% women: 31.7% Depression: total: 16.7% men:10.2% women:19.7%

The means for anxiety in the average sample and all subpopulations of ≥ 1.4 are already very close (less than 1 SD away) to the clinically significant threshold score of 1.75. This also applied to depression with means of ≥ 1.44 except for the male subpopulation. This could reflect a general disposition to quickly react with fears, worries and somatic symptoms to the unforeseen or to stress (see also: Hinton et al., 2005b).

The World Health Organization has estimated that women are 50 to 100% more likely to suffer from these disorders (WHO, 2011c) and in fact, in the present study incidence rates were much higher for women compared to men. Therefore the oversampling of female respondents has to be considered and data should be compared by gender- subpopulations where possible. Still also the scores for men were high. Compared with global figures from WHO (2011c) the point prevalence for depression found in this study is 5 times higher than the one for unipolar depression in women worldwide, although these figures are not totally comparable. The point prevalence for clinical anxiety is nearly as high as the lifelong prevalence for anxiety mentioned by WHO (2011c). In the following table prevalence rates of probable anxiety and depression from major studies in Cambodia are compared.

The present study (2011) Cambodia nationwide N= 2690 using HSCL-25		de Jong (2001/ 2003) three different Cambodian communes N=610 in Cambodia using CIDI	Dubois (2004) population from Cham province N=1320	
Anxiety	27.4%	40%	53%	
Depression	16.6%	11.5% (mood disorders)	42.4%	

Table 17: prevalence rates of anxiety and depression in comparison with other studies conducted in Cambodia

Compared with the two major previous studies conducted in Cambodia seven to ten years earlier the rates for anxiety have diminished by 30 to 50%, the ones of depression lie in between the two previous findings (see above). This could be an effect of time with an increasingly healthier second generation or due to especially high rates in the particularly badly affected province of Kampong Cham during the CW and by flooding (Dubois et al., 2004), which cannot be verified directly as this province was not randomly included in the present study. The findings by de Jong et al. (2001) measured with the CIDI (Composite International Diagnostic Interview) are closer to the findings from this study.

Studies conducted in various countries among primary care patients show that psychiatric disorders, in particular depression and anxiety, are associated with high rates of psychosocial and physical disability (Ormel et al., 1994), social deficit and loss of working capacity (Mintz et al., 1992; Sherbourne et al., 1996).

In the study by Dubois et al. (2004) in Cambodia 25.3% reported being socially impaired: regression analysis showed that psychiatric symptomatology was most strongly associated with social impairment.

IV Symptom scales and demographics

All of the symptom scales correlated significantly with sex (women more affected) and age (population > 36 more affected). Within these the anxiety scale showed the greatest difference between males and females and the C-SSI showed the greatest differences between the younger and older population. This was expected from previous statistics (WHO, 2011 c) as women are known to have a higher predisposition for worries and the C-SSI incorporates many bodily symptoms which tend to increase with age. For sex and age the HTQ-Symptoms scale showed the least differences. This can be explained via the trauma related component which makes scores more dependent of external factors (e.g. intensity and duration of traumatic experience) and more independent of intrinsic factors than other scales. Generally it can be said that gender is an important determinant of poor mental health (especially anxiety and depression) and helpseeking behavior (Oliver, Pearson, Coe & Gunnell, 2005).

Highly significant differences for marital status were observed for all four symptom scales. In all categories single respondents were least affected probably due to their lower age which has to be verified regression analysis. Widowed by respondents had the highest scores of cultural symptoms, anxiety and depression whereas divorced respondents scored highest with HTQ-Symptoms scale which may be due to the fact of re-traumatization through probable partnerviolence.

The amount of savings per respondent correlated significantly (and negatively) with all the symptom scales (-.106 with depression, -.063 with HTS, -.083 with anxiety, -.059 with C-SSI). The more savings the respondents have, the fewer symptoms they experience or vice versa the more respondents suffer from symptoms the less they can earn. The strongest correlation was found with depression. This again confirms findings from previous studies (Patel et al., 1998; Harpham, Snoxell, Grant & Rodriguez, 2005; Araya, Rojas, Fritsch, Acuna & Lewis, 2001) and the urgent need for poverty reduction as a means of prevention of mental illnesses.

The relationship between poverty and mental illnesses has also been described in a metaanalysis conducted by Patel and Kleinmann (2003). They reported an average of 20-30% prevalence rates mental of disorders developing countries. In their meta-analysis 10 out of 11 studies found a statistically significant relation poverty (especially between low education) and mental disorders.

All symptom scales correlated significantly with the educational level. Overall respondents with no or primary education have highest scores in all of the symptom scales, the highest differences were found for C-SSI and anxiety, the least for HTQ. Epidemiological data from four different lowincome and middle-income countries independently of cultural differences showed a higher vulnerability for common mental disorders for people with low education and low income than for those in upper quintiles (Patel et al., 1999). Thus relative poverty was found to be a risk factor for common mental disorders (Lund et al., 2010). However, not all studies have found a link between poverty and mental illness (Saxena et al., 2007).

Outcomes and discussion of experiences and symptoms

- Traumatic events during CW were experienced by 92.3% of all respondents aged above 36 years in high amounts as confirmed also by other studies. There seem to be differences in clusters of different experiences between provinces. Traumatic events after CW were experienced by 70.2% of respondents in comparable amounts to other studies. 86% of respondents suffered from financial worries. The majority of respondents was multiply traumatized.
- The prevalence rate of probable PTSD for the total population aged > 21 years of 2.7% found in the present survey is 7 times higher than in the general population worldwide (WHO, 2001). However it is much lower than previous findings showed. Studies using the HTQ always showed lower prevalence rates due to a high threshold score required. Changes may be due to more respondents not having experienced the CW. Women were significantly more frequently affected than men. In contrast to other studies that found high symptoms of re-experiencing the most prominent symptoms in the present study were trouble sleeping and avoidance behaviors.
- The mean for the CSSI was 1.76 out of 4.29, again with a significant gender difference (women: 1.81 versus men: 1.67). The most prominent symptoms belonged to the subscale of cultural syndromes linked to a wide spectrum of cultural beliefs, which can be regarded as a confirmation of the importance of these beliefs and associations. The rural population suffered to a significantly higher proportion of these somatic and cultural symptoms.
- The point prevalence rates with 27.4% for probable anxiety and 16.7% for probable depressive disorders were high. WHO e.g. mentions a point prevalence for unipolar depression of 1,9% for men and 3,2% for women and around 30% of the population suffering from depression, anxiety or somatic complaints. The findings described in the present study are in line and exceed respectively findings from other low-income countries. Compared with the two major previous studies conducted in Cambodia seven to ten years earlier the rates for anxiety have diminished by 30 to 50%, the ones of depression lie in between the two previous findings. This could be an effect of time with an increasingly healthier second generation or due to especially high rates in the particularly badly affected province of Kampong Cham. These facts are alarming as psychiatric disorders, in particular depression and anxiety, are associated with high rates of psychosocial and physical disability (Ormel et al., 1994), social impairment (Dubois et al., 2004) and loss of working capacity (Mintz et al., 1992; Sherbourne et al., 1996).
- All of the symptom scales correlated significantly with the sex (women more affected) and age (population > 36 more affected), marital status (singles least affected, probably due to younger age), widowed or divorced most) and the educational level (respondents with no or primary education had highest scores) and negatively with savings per capita (the more savings respondents have the less affected they are). These findings confirm findings from many other studies (Lund et al., 2010, Patel et al., 1998; Patel et al., 1999; Harpham, Snoxell, Grant & Rodriguez, 2005; Araya, Rojas, Fritsch, Acuna & Lewis, 2001) and demonstrate the need for investment in education, mental health literacy, gender balance and poverty reduction. The relationship between poverty and mental illnesses has been described already in a meta-analysis conducted by Patel and Kleinmann (2003).

V Help seeking behaviour

Alarmingly only 24.1% of the respondents reported that someone was providing help for mental health problems. In the present study most respondents seek help from a health center (including probably very small ones) or their family members. For this question multiple answers were allowed whereas a study by Dubois (2004) apparently allowed only single answers. In Dubois's study, around 30.7% sought help in the private network, 30% in public network and 12.5% in traditional medicine. The survey by MoWA found that in 14.7% of the communes there were agencies or NGOs offering general counseling or support, but only 0.9% of the inhabitants had sought help from these agencies. Here the importance of mental health literacy and the problematic issue of discrimination become visible.

Compared to official numbers of mental health services (only offered in 18 out of 967 health centers and in 9 out of 24 provinces) the numbers of services mentioned by respondents seem too high and may have been confused by respondents with lower level services.

Needs

In this study 97.3% of the respondents wished to have a psychological service at their health center and it could be concluded that 35.7% would go and see a psychologist if they had the opportunity. Knowledge of the scientific field of psychology and the true nature of the work of psychologists seems very low (qualitative results will follow).

In summary it is very obvious in this study that there is a great need for more counseling and mental health services in Cambodia. Many different issues need to be addressed in a sensitive, culturally appropriate and fitting way that utilizes existing resources (see also recommendations).

A description of experiences in the field of counseling and psychotherapy the DP has been gathering can be found in the full version of the report.

Limitations & Conclusions

- The most important limitation is the overrepresentation of women in this survey: This study was conducted during the rice planting season, which led to considerably more women (67.5%) being available for interviews in the households compared to This only became evident while examining the results. However there exists a ratio of 94.2 men to 100 women in the Cambodian population as measured by the general population Census (National Institute of Statistics, 2008) and also other studies have had this tendency (MOWA, 2009 with 57% women, de Jong, 2003 with 56.9% women).
 - The overrepresentation of women in this study was taken into account when analyzing the data and most results are presented separately for the female and male population. For future research projects it is advisable to ensure a more equal/ proportional distribution of interviews between the genders.
- Data from this survey have to be considered as only approximating the real amount of problems as many interviews (approximately 50% according to a post-study survey among the data collectors) couldn't ensure enough privacy for the respondents while answering. Data collectors tried their best to ensure privacy but often listeners would return after being sent away. This is due to a cultural Cambodian habit of curiosity and little individuality.

Traditionally there exists a lack of privacy in many circumstances and most life events are shared in public especially in rural areas. Information e.g. about diseases spreads quickly around communities. Data especially concerning the respondent's family members may be only approximating the real amount of the problems because often relatives were present during interviews or respondents may

- have feared that they would return unexpectedly and hear what they were reporting.
- Parts of the problem of holding back complete information also persist in more private interview settings. A general mistrust in unfamiliar interviewers and fears of breach of confidence or spying can be observed. This is a general problem most surveys face.
- The completion of self-assessment questionnaires depends on the respondents' honesty as it was (as in most such studies) not accompanied by a psychiatric and somatic clinical examination. Prevalence rates therefore estimate probable diagnoses, not clinical diagnoses.
- Concerning the assessment tools threshold scores used in HSCL-25 and HTQ were established on clinical populations of refugees living in the USA. Thus, interpretation of results in a community sample should be examined carefully (Dohrenwend, 1990).
- Several terms such as child abuse, torture, gambling, aggressive behavior or school problems were not further defined, so there may be a broader spectrum of interpretations that may have distorted some findings.

Conclusions

It is not astonishing that mental disorders are widespread in Cambodia considering its traumatic recent history and the adverse living conditions and poverty of its population.

However mental health services remain dwarfed, the mental health budget hasn't adequately been raised for decades and the population's needs still go largely unaddressed (McLaughlin & Wickeri, 2012). Poor mental health is proven to hinder economic development and the thriving of whole nations and many studies have stressed the economic benefit of investments in a population's mental health (WHO 2006).

- The results from the present study demonstrate a high need for more ample services for children, psychotic and addictive disorders, domestic violence, suicide prevention, anxiety, depression and PTSD.
- Much more research should be done in order to properly adjust programs and fathom understanding of causes and effects in the field of mental health in Cambodia .
- In order to face all these challenges for services and treatments a strong joint multidisciplinary effort
 is required. At the practitioner level insufficient knowledge of mutual abilities and resources and the
 existence of a professional hierarchy and lack of professional respect (with psychiatrists being of
 higher status than counselors and social workers) have thwarted cooperation and collaboration in
 many cases (Stewart, 2010).
- "The repressive nature of keeping thoughts and feelings to oneself is believed to be linked with higher incidences of mental health problems, as well as higher rates of violence. " (Stewart, 2010, p. 29). So far only a few individuals in Cambodia have acquired high level clinical and counseling skills to ensure appropriate treatment of mental disorders. Many others lack training, lack constant supervision and lack experience in the field of mental health. A tendency to overprescribe drugs in order to please the client and make up for the lack of psychosocial interventions has been reported (Stewart, 2010; McLaughlin & Wickeri, 2012).
- In response to a general discrimination and misinterpretation of mental disorders that in turn hinder the urgently needed acceptance of and access to mental health services the government should advance and support advocacy and education on mental health (WHO, 2004 b; WHO, 2004c).



	Carlson	Mollica (1993)	de Jong (2001/	Dubois (2004)	Sonis (2009)
	(1991) Cambodian refugee	Cambodian refugee population in	2003) Data from three different	population from Kampong Cham province	national sample of adults
	population in the USA n=50 using DES	Thai border refugee camps n=993 using HTQ	Cambodian communes n=610 in Cambodia using CIDI	n=1320 using Hopkins and HTQ Indochinese Version	Cambodian residents (oversampled for old age) n=1017 using PTSD Checklist
Anxiety			40%	53%	
Depression	80%	55%	11,5% (mood disorders)	42,4%	
PTSD	86%	15%	28,4%	7,3 %	11,2% 14,2 % of first generation 7,9% for younger groups
Intrusive symptoms			72.8%	47.8%	33.6%
Avoidance symptoms			59.3%	45,4%	12%
Psychosocial impairment		15-20% health impairments limiting activities		25%	mental disability for > 35: 16,0 % for < 35: 4,2 %

Table 1: research findings 1991 till 2009 of mental disorders in different samples of Cambodian population




Fig.1: Percentage of respondents having problems with family, friends or their health caused by their drinking habits (N=922)




Fig. 2: Percent of respondents having problems with family or friends caused by their gambling (N=143)




Fig. 3: HTQ-Symptoms experienced by respondents with means > 1.43 (N = 2681-2689)

- The symptom with the highest mean score was trouble sleeping followed by feeling irritable, avoidance of activities and thoughts associated with traumatic experiences and difficulty performing work or daily tasks
- Feelings of emotional numbing (M = 1.21), shame (M = 1.28) and dissociative experiences (loss of memory (M = 1.29), feeling detached (M = 1.28), feeling split (M = 1.23)) were least experienced




Fig. 4 : CSSI items experienced by respondents with a mean > 1.76 (N = 2683-2689)

- The most prominent item from the CSSI was the syndrome of `thinking too much', followed by the syndrome of `numb arms and legs' and the symptom of `standing and feeling dizzy'
- `Sleep paralysis' (M = 1.16), `poor appetite' (M = 1.43) and `cold hands and feet' (M = 1.48) were the least frequent items




Fig. 5: most prominent HSCL 25 anxiety symptoms with mean > 1.51 (N = 2687-2689)





Fig. 6: HSCL-25 symptoms of depression with means > 1.45 (N = 2686-2689)



Almqvist K., Broberg, A. (2003): Young children traumatized by organized violence together with their mothers - The critical effects of damaged internal representations. In: *RAHD* 5 (4), P. 367–380.

Araya R,. Rojas G., Fritsch R., Acuna J., Lewis G. (2001): Common mental disorders in Santiago, Chile: Prevalence and socio-demographic correlates. In: *Brit-ish Journal of Psychiatry* (178), P. 228–233.

Bar-On J., Eland R. J., Klebe, D. (1998): Multigenerational Perspectives on Coping with the Holocaust Experience: An Attachment Perspective for Understanding the Developmental Sequelae of Trauma across Generations. In: *International Journal of Behavioral Development* 22 (2), P. 315–338.

Berthold S.M., Gray G. (2011): Post-Traumatic Stress Reactions And Secondary Trauma Effects At Tribunals: The ECCC Example. In: Reicherter, Youk Chhang, van Schaack (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Og The Khmer Rouge. Phnom Penh: DCCAM (Documentation Series, No 17), P. 92–115.

Bertrand D. (2005): The therapeutic role of Khmer mediums (kru boramei) in contemporary Cambodia. In: *Mental Health, Religion & Culture*.

Bit S. (1991): The Warrior Heritage: A psychological perspective of Cambodian trauma. El Cerrito, CA: Seanglim Bit (self-published). In: *El Cerrito, CA: Seanglim Bit (self-published)*.

Boehnlein J.K., Kinzie J.D. (2011): The Effect Of The Khmer Rouge On The Mental Health Of Cambodia And Cambodians. In: Reicherter, Youk Chhang, van Schaack (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Of The Khmer Rouge. Phnom Penh: DCCAM (Documentation Series, No 17), P. 33–41.

Bremner J.D., Southwick S.M., Darnell A., Charney D.S. (1996): Chronic PTSD in Vietnam combat veterans: course of illness and substance abuse. In: *American Journal of Psychiatry* (153), P. 369–375.

Brewin C.R., Dalgeish T., Joseph S. (1996): A dual representation theory of posttraumatic stress disorder. Psychological Review (103), P. 670–686.

Bride B. E. (2004): Development and Validation of the Secondary Traumatic Stress Scale. In: *Research on Social Work Practice* 14 (1), P. 27–35.

Briere J., Scott C., Wheaters F. (2005): Peritraumatic and Persistence Dissociation in the presumed etiology of PTSD. In: American Journal of Psychiatry 162 (12), P. 2295–2301.

Briere J., Scott C. (2006): Principles Of Trauma Therpy: A Guide To Symptom, Evaluation, and Treatment. Thousand Oaks, CA: Sage Publications.

Briere J., Hodges M., Godbout N. (2010): Traumatic Stress, Affect Dysregulation, and Dysfunctional Avoidance: A Structural Equation Model. In: *Journal of traumatic stress* 23 (6), P. 767–774.

Bromet E., Sonnega A., Kessler R.C. (1998): Risk factors for DSM-III-R posttraumatic stress disorder: findings from the National Comorbidity Survey. In: *American Journal of Epidemiology* (15), P. 353–361.

Bunthoeun P. (1998): Explanatory models of mental health problems of outpatients at the psychiatric department of Preah Norodom Sihanouk Hospital. Postgraduate thesis, Phnom Penh, Cambodia. Cambodian Mental Health Training Program.

Calless I.T., Machleidt W., Ziegenbein M., Haltenhof H. (2007): Suizidalität im Kulturvergleich. Transcultural Aspects of suicidal behavior. In: *Fortschritt Neurologische Psychiatrie* 75, P. 653–664.

Cambodian Daily (March 23rd 2011): Teen Students Planning or Attempting Suicide, P. 22.

Cambodian Daily (March 10th 2011): Suicide Death Toll 542 in 2010, Up 35 Percent, P.23.

Carlson E.B., Rosser-Hogan R. (1991): Trauma experiences, posttraumatic stress, dissociation, and depression in Cambodian refugees. In: *American Journal of Psychiatry* (148), P. 1548–1551.

Centers for Disease Control and Prevention (CDC). (2009): Suicide: facts at a glance. Edit. v. Statistics Query and Reporting System (WISQARS). Centers for Disease Control and Prevention (CDC). http://www.cdc.gov/violenceprevention/pdf/suicidedatasheet-a.pdf retrieved 28.10.2011.

Chinese Women's Research Network Xinhua: UN Chief Says "No Health without Mental Health" on International Day. http://en.wsic.ac.cn/academicnews/3396.htm retrieved 02.11.2011.

Ciorciari J.D., Heindel A. (2011): Trauma In The Courtroom. In: Reicherter Youk Chhang van Schaack (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Of The Khmer Rouge. Phnom Penh: DCCAM (Documentation Series, No 17), P. 122–138.

Clark D.M., Ehlers A. (2004): Posttraumatic stress disorder: From cognitive theory to therapy. In: R.L Leahy (Edit.): Contemporary cognitive therapy. New York: Guilford Press.

Cohen J.A., Mannarino A.P. (2006): Treating Trauma and Traumatic Grief in Children and Adolescents. New York: Guilford Press.

Crookes A. (March 17th 2011): Substance Abuse Workshop. Phnom Penh, March 17th 2011.

Davidson J.R.T., Foa E.B. (Edit.) (1993): Posttraumatic Stress Disorder: DSM-IV and Beyond. Washington DC: American Psychiatric Press.

De Jong J. T.V.M., Komproe I., de Ommeren M. (2003): Common mental disorders in postconflict settings. In: *The Lancet* 361 (9375), P. 2128,

De Jong J.T., Komproe T.V.M., Ivan H., Van Ommeren M., El Masri M., Araya M. et al. (2001): Lifetime events and posttraumatic stress disorder in 4 postconflict settings. In: *Journal of the American Medical Association* (285), P. 555–562.

De Jong, J. (Edit.) (2002): Trauma, War and Violence: Public Mental Health in Socio-Cultural Context. New York: Kluwer Academic/Plenum Publishers.

Desjarlais R., Eisenberg L., Good B., Kleinman A. (1995): World mental health. problems and priorities in low income countries. Oxford, UK: Oxford University Press.

Dicklitch S., Malik A. (2010): Justice, Human Rights, and Reconciliation in Postconflict Cambodia. In: *Human Rights Review* 11 (4), P. 515–530.

Dohrenwend B.P (Edit.) (1998): Adversity, Stress and Psychopathology. New York: Oxford University Press.

Dubois V., Tonglet R., Hoyois P., Sunbaunat K., Roussaux J., Hauff E. (2004): Household Survey of Psychiatric Morbidity in Cambodia. In: *International Journal of Social Psychiatry* 50 (2), P. 174–185.

Dy K. (2009) A History of Democratic Kampuchea, DCCAM in partnership with the Cambodian Ministry of Education.

Eisenbruch M., de Jong J.T.V.M, van de Put W. (2004): Briniging order out of chaos: A culturally competent approach to managing the problems of refugees and victims of organized violence. In: *Journal of traumatic stress* 17 (2), P. 123–135.

Eungprabhant V. (1975): Suicide in Thailand 60-70. In: *Forensic Science* 5, P. 43–51.

Fals-Stewart W. (2003): The occurrence of partner physical aggression on days of alcohol consumption. In: *Journal of Consulting and Clinical Psychology* (71), P. 41–52.

Ferrada-Noli M., Asberg M., Ormstad K., Lundin T., Sundbom E. (1998): Suicidal Behavior After Severe Trauma: Part 1 PTSD Diagnosis, Psychiatric Comorbidity, and Assessments of Suicidal Behavior. In: *Journal of traumatic stress* 1 (11). P.103-112

Field N. P., Chhim S. (2008): Desire for Revenge and Attitudes Toward the Khmer Rouge Tribunal Among Cambodians.. Journal of Loss and Trauma, 352-372. In: *Journal of Loss and Trauma*, P. 352–372.

Field N.P. (2011): Intergenerational Transmission Of Trauma Stemming From The Khmer Rouge Regime: An Attachment Perspective. In: Reicherter Youk Chhang van Schaack (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Of The Khmer Rouge. Phnom Penh: DCCAM (Documentation Series, No 17), P. 70–85.

Foa E., Rothbaum O. (1998): Treating the trauma of rape. New York: Guilford Press.

Follette V.M., Polusny M., Brechtle A.E., Naugle A.E. (1996): Cumulative trauma: the impact of child sexual abuse, adult sexual assault, and spouse abuse. In: *Journal of traumatic stress* (9), P. 25–35.

Fox A. (2008): Sociocultural Factors that Foster or Inhibit Alcohol-related Violence. In: Alcohol and Violence: Exploring Patterns and Responses, International Center for Alcohol Policies (ICAP)

Green B.L (1990): Defining trauma: terminology and generic stressor dimensions. In: *Journal of applied social psychology* (20), P. 1632–1642.

Green B. L., Goodman L.A., Krupnik J.L., Corcoran C.B., Petty R.M., Stockton P., Stern N.M. (2000): Outcome of single versus multiple trauma exposure in a screening sample. In: *Journal of traumatic stress* (13), P. 271–286.

GTZ (2009): Violence Against Women: 2009 Follow-Up Survey & Final Study Report. GIZ. Phnom Penh, http://www.un.org.kh/undp/knowledge/publications/violence-against-women-2009-follow-up-survey.

Harpham T., Snoxwell S., Grant E., Rodriguez C. (2005). Common mental disorders in a young urban population in Colombia. In: *British Journal of Psychiatry* (187), P. 161–167.

Harris R. (2009): ACT made simple: An Easy-to-Read Primer on Acceptance and Commitment Therapy. Oakland: New Harbinger Publications.

Hasin D., Stinson F. S., Ogburn E., Grant B. (2007): Prevalence, Correlates, Disability, and Comorbidity of DSM-IV Alcohol Abuse and Dependence in the United States Results From the National Epidemiologic Survey on Alcohol and Related Conditions. In: *Archives on general psychiatry* (64 (7)), P. 830–842.

helpguide.org: Parenting bonding reactive attachment disorder. http://helpguide.org/mental/parenting_ bonding_reactive_attachment_disorder.htm retrieved 28.10.2011.

Henderson D.C., Mollica R.F., Tor S., Lavelle J., Culhane M.A., Hayden D. (2005): Building Primary Care Practitioners' Attitudes and Confidence in Mental Health Skills in a Post-Conflict Society: A Cambodian Example. In: *Journal of Nervous & Mental Disease* (193 (3)), P. 551–559.

Herbst P. (1992): From helpless victim to empowered survivor: Oral history as a treatment for survivors of torture. In: *Woman and Therapy* (13(1/2)), P. 141–154.

Herlihy J., Scragg P., Turner S. (2002): Discrepancies in autobiographical memories— implications for the assessment of asylum seekers: repeated interviews study. In: *BMJ* 324, P.324–327.

Hinton D., Um K., Ba P. (2001): Kyol Goeu ('Wind Overload') Part I: A Cultural Syndrome of Orthostatic Panic among Khmer Refugees. In: *Transcultural Psychiatry* 38 (4), P. 403–432.

Hinton D., Hinton S., Um K., Chea A., Sak S. (2002): The Khmer 'Weak Heart' Syndrome: Fear of Death from Palpitations. In: *Transcultural Psychiatry* 39 (3), P. 323–344.

Hinton D. E., Chhean D., Pich V., Safren S. A., Hofmann S. G., Pollack, M. (2005): A randomized controlled trial of cognitive-behavior therapy for Cambodian refugees with treatment-resistant PTSD and panic attacks: A cross-over design. In: *J. Traum. Stress* 18 (6), P. 617-629.

Hinton D. E., Pich V., Chhean D. (2005a): 'The Ghost Pushes You Down': Sleep Paralysis-Type Panic Attacks in a Khmer Refugee Population. In: *Transcultural Psychiatry* 42 (1), P. 46–77.

Hinton D. E., Chhean D., Pich V., Safren S. A., Hofmann, S., Pollack M. (2005b): A randomized controlled trial of cognitive-behavior therapy for Cambodian refugees with treatment-resistant PTSD and panic attacks: A cross-over design. In: *J. Traum. Stress* 18 (6), P. 617–629.

Hinton D. E., Pich, V., Marques L., Nickerson A., Pollack M. (2010): Khyâl Attacks: A Key Idiom of Distress Among Traumatized Cambodia Refugees. In: *Culture, Medicine, and Psychiatry* 34 (2), P. 244–278.

Hinton D. E., Hinton A., Eng K-Th., Choung S. (2011): PTSD Severity And Key Idioms Of Distress Among Rural Cambodians: The Result Of A Needs Assessment Survey. In: Reicherter, Youk Chhang, van Schaack (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Of The Khmer Rouge. Phnom Penh: DCCAM (Documentation Series, No 17), P. 47–66.

Hinton D. E., Lewis-Fernandez R. (2011): The cross-cultural validity of posttraumatic stress disorder: Implications for DSM-5. In: *Depression and Anxiety* (28), P. 783–801.

Hinton D.E., Hinton A. L., Chhangd Y., Eng K-Th., Choungd S. (2012): PTSD and Key Somatic Complaints and Cultural Syndromes among Rural Cambodians: The Results of a Needs Assessment Survey. In: *Medical Anthropology Quarterly* 26 (3),P- 383-407.

Hoshmand L. T. (2006): Culture-informed theory, reserach and practise in global trauma work. In: *Journal of Counselling* 13 (1), P. 145–160.

Jacobsen T. (2006): Punishing the perpetrators of the Cambodian Genocide. In: *Asia Insights* (3), P. 6–7.

Jegannathan B., Kullgren G. (2011): Gender differences in suicidal expressions and their determinants among young people in Cambodia, a post-conflict country*. In: *BMC Psychiatry 2011, 11:47* 11 (47).

Johnson L. R., Mayanja M.K., Bangirana P., Kizito S. (2009): Contrasting Concepts of Depression in Uganda: Implications for Service Delivery in a Multicultural Context. In: *American Journal of Orthopsychiatry* (79 (2)), P. 275–289.

Kessler R.C., Sonnega A., Bromet E., Nelson C.B. (1995): Posttraumatic stress disorder in the National Comorbidity Survey. In: *Archives on general psychiatry* (52), P. 1048–1060.

Khann S. (2010): Domestic Violence Against Women in Cambodia:An Examination of Women's Marital Resources, Husband's Control and Family Structure. Royal University of Phnom Penh, Phnom Penh. Faculty of Social Sciences and Humanities Department of Psychology.

Killpatrick D.G., Resnick H.S (1993): PTSD associated with exposure to criminal victimization in clinical and community populations. In: J.R.T Foa E.B Davidson (Edit.): Posttraumatic Stress Disorder: DSM-IV and Beyond.; Washington, DC: American Psychiatric Press, P. 113-143.

Killpatrick D.G., Resnick H.S., Saunders B.E., Best C.L (1998): Rape, other violence against women, and posttraumatic stress disorder: critical issues in assessing the adversity-stress-psychopathology relationship. In: B.P Dohrenwend (Edit.): Adversity, Stress and Psychopathology. New York: Oxford University Press, P. 161–176.

Kingdom of Cambodia, Law on Prevention of Domestic Violence and Protection of the Victims (2005) http://www.bno.com.kh/Cambodia%20Laws/205_LAW_Prevention%20of%20Domestic%20Violence,Protection%20of%20Victims,%20Eng.pdf retrieved 13.11.2011

Kirchenbauer N., Burchert S., Taing S., Bockers E., Knaevelsrud C. (2010): The Survivors' Voices: Attitudes on the ECCC, the Former Khmer Rouge and Experiences with Civil Party Participation, Berlin, Germany. Behandlungszentrum für Folteropfer e.V. (Center for the Treatment of Torture Victims).

Lambert J. (1986). Khmer refugees in *Dallas: Medical decisions in the context of pluralism.* PhD dissertation, Southern Methodist University, UMI Dissertation Service, Ann Arbor, MI.

Leahy R.L (Edit.) (2004): Contemporary cognitive therapy. New York: Guilford Press.

Leo D., Milner A., Xiandong W. (2009): Suicidal Behavior in the Western Pacific Region Characteristics and Trends. In: *Suicide and Life Threatening Behavior* 39 (1), P. 70–81.

Lotrakul M. (2006): Suicide in Thailand during the period 1998-2003. In: *Psychiatry and Clinical Neuroscience* (60), P. 90–95.

Lund C., Breen A., Flisher A.J., Kakuma R., Corrigall J., Joska J.A., Swartz L., Patel V. (2010): Poverty and common mental disorders in low and middle income countries: A systematic review. In: Soc Sci Med. 71(3), P. 517-28. Epub 2010 May 12.

Marshall G.N., Elliott M. N., Schell T. L. (2009): Prevalence and correlates of lifetime disordered gambling in Cambodian refugees residing in Long Beach, CA. In: *Journal of Immigrant Minority Health* (11(1)), P. 35–40.

Marshall G.N. (2009): Prevalence and Correlates of Lifetime Disordered Gambling in Cambodian refugees Residing in Longbeach CA. In: *Immigrant & Minority Health* (35).

Mathers C.D., Loncar D. (2005): Updated projects of global mortality and global burden of disease, 2020: Global Burden of Disease Study. In: *The Lancet*, P. 1498–1504.

Mc Cay M., Wood J.C., Brantley J. (2007): The dialectical behavior therapy skills workbook- practical DBT Exercises for Learning Mindfulness, interpersonal Effectiveness, Emotion Regulation & Distress Tolerance. Oakland: New Harbinger Publications.

McDaid D., Knapp M., Raja S. (2008): Barriers in the mind: promoting an economic case for mental health in low- and middle- income countries. In: *World Psychiatry*, P. 79–86.

McLaughlin D., Wickeri E. (2012): Special Report: Mental Health and Human Rights in Cambodia. In: Fordham International Law Journal (35), P. 895-967

Miles G., Miles G., (2006a), Stop Violence against Us! Summary Report I,. Tearfund. http://www.globalempowerment.org/policyadvocacy/pahome 2.5.nsf allreports/959D0DDEB18DA9668825714D002 50470/\$file/StopViolenceAgainstUs!1_web.pdf. retrieved 01.11.2011.

Miles G. (2006b): Stop Violence Against Us! Summary Report 2, A Preliminary National Research Study into the Prevalence & Perceptions of Cambodian Children to Violence Against and By Children in Cambodia. http://www.globalempowerment.org/policyadvocacy/pahome2.5.nsf allreports/6CA8A2FBB1685F8B8825714D007D4E50/\$file/Stop%20Violence%20against%20Us!2_web.pdf retrieved 01.11.2011.

Ministry of Health (2003): National Mental health Plan 2003-2022, Final Draft.

Ministry of Health (2007): Cambodia Health Information System: Review and Assessment. Department of Planning and Health Information. Phnom Penh.

Ministry of Health (2011): Mental Health and Substance Misuse. Strategy Plan 2011-2015. National Program for Mental Health, World Health Organization. Phnom Penh, Cambodia.

Ministry of Labour and Vocational Training (2011): National social security fund, http://www.nssf.gov.kh/.

Ministry of Women's Affairs (2008). A fair share for women: Cambodia gender assessment 2008. Phnom Penh: Cambodia.

Mintz J., Mintz L.I., Arruda M.J., Hwang S. (1992): Treatment of depression and the functional capacity to work. In: *Archives on general psychiatry* (49), P. 761–781.

Mollica R.F., Wyshak G., Marneffe D., Khuon F., de Lavelle J. (1987): Indochinese versions of the Hopkins Symptom Checklist-25: a screening instrument for the psychiatric care of refugees. In: *American Journal of Psychiatry* (144 (4)), P. 497–500.

Mollica R. F., Donelan K., Tor S., Lavalle J., Elias C., Frankel M., Blendon R.J. (1993): The effect of trauma and confinement on functional health and mental health status of Cambodians living in Thailand- Cambodia border camps. In: *JAMA* (270), P. 581–586.

Mollica R.F., Caspi-Yavin Y., Lavelle J., Tor S., Yang T., Chan S., Pham T., Ryan A., Marneffe D. (1996): The Harvard Trauma (HTQ) Manual: Cambodian, Laotian, and Vietnamese versions. In: *Torture* (Supll.1), P. 19–42.

Mollica R. F., McInnes K., Pham T., Smith Fawzi M.C., Murphy E., Lin L. (1998a): The dose-effect relationships between torture and psychiatric symptoms in Vietnamese ex-political detainees and a comparison group. In: *Journal of Nervous & Mental Disease* (186), P. 543–553.

Mollica R., McInnes K., Poole C., Tor S. (1998b): Doseeffect relationships of trauma to symptoms of depression and post-traumatic stress disorder among Cambodian survivors of mass violence. In: *British Journal of Psychiatry* (173), P. 482–488.

Mollica R.F., McDonald L.S., Massagli M.P., Silove D.M. (2004): Measuring Trauma, Measuring Torture; Instructions and Guidance on the utilization of the Harvard Program in Refugee Trauma's Versions of The Hopkins Symptom Checklist-25 (HSCL-25) & The Harvard Trauma Questionnaire (HTQ), © Harvard Program in Refugee Trauma

Munyas B. (2008): Genocide in the Minds of Cambodian Youths: Transmitting (Hi)stories of Genocide to Second and Third Generations in Cambodia. In: *Journal of genocide 10 (3,) P. 413-439.*

Murray C.J., Lopez A. D. (1997): Alternative projections of mortality and disability by cause 1990-2002-2030: data sources, methods and results. Edit. v. Geneva: World Health Organization.

National Institute of Public Health, National Institute of Statistics and ORC Macro (2006): Cambodia demographic and health survey 2005. National Institute of Public Health, National Institute of Statistics and ORC Macro. Phnom Penh, Cambodia, and Calverton, Maryland. USA.

National Institute of Statistics (2008): Kingdom of Cambodia: General Population Census of Cambodia 2008. Ministry of Planning. Phnom Penh, Cambodia.

Nelson B.D., Fernandez W.G., Galea S., Sisco S., Dierberg K., Gorgieva G.S., Nandi A.K., Ahern J., Mitrovic M., Van Rooyen M., Vlahov D. (2004): Warrelated psychological sequelae among emergency department patients in the former Republic of Yugoslavia. In: *BMC Med* 2 (1), P. 22.

Nickerson A. Hinton D. E. (2011): Anger Regulation in Traumatized Cambodian Refugees: The Perspectives of Buddhist Monks. In: *Cult Med Psychiatry* 35 (3), P. 396–416.

Ohayon M., Shapiro C. (2000): Sleep disturbances and psychiatric disorders associated with posttraumatic stress disorder in the general population. In: *Comprehensive Psychiatry* (41), P. 469–478.

Oliver M.I., Pearson N., Coe N., Gunnel D. (2005): Help-seeking behaviour in men and women with common mental health problems: cross-sectional study. In: *British Journal of Psychiatry* (186), P. 297–301.

Ormel J, VonKorff M, Ustun TB, Pini S, Korten A, Oldehinkel T (1994): Common mental disorders and disability across cultures: results from the WHO collaborative study on psychological problems in general health care. Journal of the American Medical Association. In: *JAMA* 272, P. 1741–1748.

Otto M. W., Deckersback T. (1998): Cognitive-behavioral therapy for panic disorder: Theory, strategies, and outcome. In: J. Pollack M. Rosenbaum (Edit.): Panic disorder and its treatment. New York: Marcel Dekker, P. 181–205.

Pact Cambodia (2004): Promoting decentralization through partnership. Edit. v. Pact Cambodia. http://www.pactcambodia.org/Publications/Decentralization/Partnership_Handbook_English.pdf last retrieved 02.011.2011.

Paradis C., Friedman S., Hatch M. (1997): Isolated sleep paralysis in African Americans with panic disorder. In: *Cultural Diversity and Mental Health* (3), P. 69 –73.

Patel V., Pereira J., Coutinho L., Fernandes R., Fernandes J., Mann A. (1998): Poverty, psychological disorder and disability in primary care, attenders in Goa, India. In: *British Journal of Psychiatry* (172), P. 533–566.

Patel V., Araya R., Lima M., Ludermir A., Todd C. (1999): Women, poverty and common mental disorders in four restructuring societies. Soc Sci Med. In: *Social Science and Medicine* (49), P. 1461–1471.

Patel V., Kleinmann A. (2003): Poverty and common mental disorders in developing countries, Bulletin of the World Health Organization 2003, 81 (8)

Patel V., Araya R., Chatterjee S. (2007): Treatment and prevention of mental disorders in low and middle income countries. In: *Lancet* (370),P. 991–1005

Phan, T., Silove D. (1999): Overview of Indigenous Descriptions of Mental Phenomena and the Range of Traditional Healing Practices Amongst the Vietnamese. In: *Transcultural Psychiatry* (36(1)), P. 79–94.

Pham P., Vinck P., Balthazard M., Hean S. (Edit.) (2011): AFTER THE FIRST TRAIL. A Population-Based Survey on Knowldege and Perception Of Justice And The Extraordinary Chambers In The Courts Of Cambodia.

Pickwell, S.M (1999): Health of Cambodian Refugees. In: *Journal of Immigrant Health* 1 (1), P. 49–52.

Pinheiro P.S. (2006): World Report on Violence against Children. Edit. v. United Nations Secretary-General. Geneva.

Prince M., Patel V., Saxena S., Maj M., Maselko J., Phillips M.R., Rahman A.: No health without mental health. In: *The Lancet* (370 (9590)), P. 859–877.

Rechtman R. (2000): Stories of Trauma and Idioms of Distress: From Cultural Narratives to Clinical Assessment Transcultural Psychiatry, P. 403–415.

Rechtman R. (2006): The Survivor's Paradox: Psychological Consequences of the Khmer Rouge Rhetoric of Extermination. In: *Anthropology & Medicine*, P. 1–11.

Reicherter D., Boehnlein J., Stewart J. (2011): Analysis Of Trauma-Related Mental Health Resources In Cambodia: Consensus Ideas For An Improved Method. In: Reicherter D., Youk C., van Schaak B.(Edit.): Cambodia's Hidden Scars: Trauma Psychology In The Wake Of The Khmer Rouge. DCCAM, Documentation Series No 17, P. 206–217.

Reicherter D., Aylward A. (2011): The Impact Of War And Genocide On Psychiatry And Social Psychology. In: Reicherter D., Youk C., van Schaak B. (Edit.): Cambodia's Hidden Scars: Trauma Psychology In The Wake Of The Khmer Rouge. DCCAM, Documentation Series No 17, P. 14-31.

Reicherter D., Eng K-T (2011): Cambodian Perspectives on The Resources For Trauma Related Mental Health. In: Reicherter D., Youk C., van Schaak B. (Edit.): Cambodia's Hidden Scares. Trauma Psychology In The Wake Og The Khmer Rouge. Phnom Penh: DCCAM, Documentation Series, No 17, P. 191–204.

Rifà H., Franco B. (2011). Violence Against Women in Partner Relationship. Indigenous Peoples in Ratanakiri Province: A survey study. Paper presented at the Workshop on Domestic Violence among Indigenous Communities, Phnom Penh: Psicólogos Sin Fronteras ONGD, in press

Rosenbaum J., Pollack M. (Edit.) (1998): Panic disorder and its treatment. New York: Marcel Dekker.

Rousseau C., Drapeau A., Rahimi S. (2003): The complexity of trauma response: a 4-year follow-up of adolescent Cambodian refugees. In: *Child Abuse & Neglect* 27 (11), P. 1277–1290.

Saxena S., Sharan P., Garrido M., Saraceno B. (2006): World Health Organization's Mental Health Atlas 2005: Implications For Policy Development. In: *World Psychiatry* (5), P. 179–184.

Saxena S., Thornicroft G., Knapp M., Whiteford H. (2007): Resources for mental health: scarcity, inequity, and inefficiency. In: *The Lancet* 370 (9590), P. 878–889

Schaala S., Elberta T., Neunerb F. (2009): Narrative Exposure Therapy versus Interpersonal Psychotherapy A Pilot Randomized Controlled Trial with Rwandan Genocide Orphans. In: *Psychotherapy and Psychosomatics* (78), P. 298–306.

Seoung S. (2011): Psychological Disaster Management in Cambodia - a study of a short term intervention using the Emergency Room Protocol with victim from the 2010 Koh Pich stampede, at Preah Kosamak Hospital. Masterthesis. Royal University of Phnom Penh, Phnom Penh. Department of Psychology.

Shankar B., Saravanan B., Jacob K.S. (2006): Explanatory Models of Common Mental Disorders among Traditional Healers and Their Patients in Rural South India. In: *International Journal of Social Psychiatry* (52 (3)), P. 221–233.

Sherbone C.D., Wells K.B., Judd L.L (1996): Functioning and well-being of patients with panic disorder. In: *American Journal of Psychiatry* (153), P. 213–218.

Somasundaram D.J., Sivayokan S. (1994): War trauma in a civilian population. In: *British Journal of Psychiatry* (165), P. 524–527.

Somasundaram D.J., van de Put W.A.C. (1999): Mental Health Care in Cambodia. In: *Bulletin of the World Health Organization* 77 ((3)), P. 275–277.

Sonis J., Gibson J.L, .de Jong J.T.V.M., Field N.P., Hean S., Komproe I. (2009): Probable Postraumatic Stress Disorder and Disability in Cambodia. Associations with perceived Justice, Desire for Revenge and Attitudes towards the khmer rouge trials. In: *JAMA* 302 (5), P. 527–536.

Stewart J., Tsong Y., Peou P.C. (2010): Mental Health in Cambodia: A qualitative Evaluation. In: *unpublished*.

Stewart S. H. (1996): Alcohol Abuse in Individuals Exposed to Trauma: A Critical Review. Psychological Bulletin (No. 1). Online verfügbar unter http://my.ilstu.edu/~dfgrayb/Personal/alcohol%20and% 20trauma.pdf, zuletzt geprüft am 28.10.2011.

Stockwell A., Whiteford H., Townsend C., Stewart D. (2005): Mental health policy development: case study of Cambodia. In: *Australasian Psychiatry* 13 (2), P. 190–194.

Thornicroft G., Tansella M. (2004): Components of a modern mental health service: a pragmatic balance of community and hospital care:overview of systematic evidence. In: *British Journal of Psychiatry* (185), P. 283–290.

United Nations Development Program (2008): Human Development Report 2007–2008. http://hdr.undp. org/en/statistics/. Retrieved 29.10.2011

United Nations Development Program (2010): Human Development Report. Human Development Index. http://hdrstats.undp.org/en/countries/profiles/KHM. http://hdrstats.und

United Nations World Drug Report (2010): UNDP Annual Report 2010 Cambodia, empowered lives, resilient nations.

Van Dijk J.A., Schoutrop M.J., Spinhoven P. (2003): Testimony therapy: treatment method for traumatized victims of organized violence. In: *Journal of Psychotherapy* (57(3)), P. 361–373.

Van de Put W., Eisenbruch M. (2002): The Cambodian Experience. In: de Jong J. (Edit.): Trauma, War and Violence: Public Mental Health in Socio-Cultural Context. New York: Kluwer Academic/Plenum Publishers, P. 93–156.

Van de Put W., van der Veer G. (2005): Counselling in Cambodia: cultural competence and contextual costs. In: *Intervention*, P. 87–96.

Van Schaak B., Reicherter D., Youk C. (Edit.) (2011): Cambodia's Hidden Scars: Trauma Psychology In The Wake Of The Khmer Rouge. Documentation Center of Cambodia (www.dccam.org). Documentation Series No 17.

Van Steel Z., Chey T., Silove D., Marnane C., Bryant R. A., Ommeren M. (2009): Association of Torture and Other Potentially Traumatic Events With Mental Health Outcomes Among Populations Exposed to Mass Conflict and Displacement A Systematic Review and Meta-analysis. In: *JAMA* 302 (5), P. 537–549.

Wagner R.S., Sergio A. B., Cleusa F. P., Martin P. (2009): Exposure to violence and mental health problems in low and middle-income countries: a literature review. In: *Revista Brasileira de Psiquiatria* (31), P. 49 –57.

Weissman M.M., Bland R.C., Canino G.J., Greenwald S., Hwu H.G., Joyce P.R., Karam E.G., Lee C.K., Lellouch J., Lepine J.P., Newman S.C., Rubio-Stipec M., Wells J.E., Wickramaratne P.J., Wittchen H.U., Yeh E.K (1999): Prevalence of suicide ideation and suicide attempts in nine countries. In: *Psychological Medicine* (29(1)), P. 1–17.

Wiley-Exley E. (2007): Evaluations of community mental health care in lowand middle-income countries: A 10-year review of the literature. In: *Social Science & Medicine* (46), P. 1231–1241.

Woods L.N (2008): Sound the Alarm: Reporting violence against children in Cambodia. Unicef. http://www.scribd.com/doc/30801758/SOUND-the-ALARM-Reporting-Violence-Against-Children-in-Cambodia.

World Bank (1994): World development report 1993: investing in health. New York: Oxord University Press.

World Health Organization (1990): The introduction of a mental health component into primary health care. Edit. v. World Health Organization. Geneva, Switzerland

World Health Organization (1996): Mental health of refugees-Manual. World Health Organization, Geneva, Switzerland. http://helid.digicollection.org/en/d/Jh0223e/2.2.html, retrieved 07.10.2011.

World Health Organization (2001): The World Health Report 2001: Mental Health: New understanding, new hope. http://www.who.int/whr/2001/en/whr01 en.pdf retrieved 14.11.2011

World Health Organization (2004a): World Mental Health Consortium. Prevalence, severity and unmet need for treatment of mental disorders in the World Mental Health Organization world mental health surveys. In: *Journal of the American Medical Association* (291), P. 2581–2590.

World Health Organization (2004b): Prevention of Mental Disorders Organization, Effective interventions and policy options summary report. Edit. v. World Health Organization. Department of Mental Health and Substance Abuse, Prevention Research Centre of the Universities of Nijmegen.

World Health Organization. (2004c): Promoting mental health: concepts, emerging evidence, practice: summary report. Edit. v. World Health Organization. Geneva, Switzerland.

World Health Organization (2008): Mental Health Gap Action Programme: scaling up care for mental, neurological and substance use disorders. World Health Organization. Geneva, Switzerland

World Health Organization (2011a): Suicide prevention. http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/ retrieved 27.10.2011.

World Health Organization (2011b): Suicide rates. http://www.who.int/mental_health/prevention/suicide/suiciderates/en/retrieved 27.10.2011.

World Health Organization (2011c): Gender and women's mental health. http://www.who.int/mental_health/prevention/genderwomen/en/retrieved 28.10.2011.





